



# **BACTICID AF**

## Alcohol based rapid disinfectant for medical devices

**Bacticid AF** is alcohol-based rapid disinfectant for non-invasive medical devices and alcohol resistant surfaces. Perfect product in higher hygiene requirements, where rapid and regular surface disinfection is required. The product comes as a ready solution which is very convenient to use. It does not contain aldehydes, fragrance and colorants and is suitable for all alcohol-resistant surfaces.

**Bacticid AF** has a broad spectrum of bactericidal (incl. MRSA, VRE), mycobactericidal (incl. Mycobacterium avium and Mycobacterium terrae), limited virucidal (incl. HBV, HIV, HCV, Adeno, Noro, Vaccinia), fungicidal and yeasticidal activity.

#### **APPLICATION AREAS:**

Alcohol-based rapid disinfection of medical devices and other surfaces, where short contact times are required, for example in the following areas:

- Hospitals and other medical institutions
- Operating theatres and procedure rooms
- Intensive care unit
- In accident and emergency departments
- Dialysis departments
- Family doctor offices, first-aid stations
- Ambulances
- Laboratories and diagnostics.
- Dentist's offices
- For home care of patient
- Food industry and food processing areas
- Pharmaceutical and cosmetics industry

#### **USAGE:**

BACTICID AF is ready for use solution and can be applied on the surfaces by wiping or spraying.

Surface disinfection by wiping: Apply a sufficient amount of BACTICID AF to a suitable single-use non-woven wipe and wipe the surfaces.

Surface disinfection by spraying: Spray BACTICID AF onto surfaces from a distance of at least 30 cm. Recommended use for difficult to reach surfaces.







Ensure complete coverage. The surfaces must remain visibly moistened for 30 seconds to take effect. Maximum application rate 50 ml/m<sup>2</sup>.

Bacticid AF is an effective product for rapid disinfection of surfaces like operating tables, examination couches, stretchers, patient chairs, wheelchairs, medical equipment, trays, trolleys, hospital and maternity beds.

If the surface tolerance is unknown, try it out on an inconspicuous area first. This product is not suitable for acrylic or Plexiglas® nor alcohol soluble varnish.

### MICROBIOLOGY:

Efficacy	Concentration	Contact time
Escherichia coli	Ready to use	30 sec
Pseudomonas aeruginosa	Ready to use	30 sec
Staphylococcus aureus	Ready to use	30 sec
Enterococcus hirae	Ready to use	30 sec
Staphylococcus aureus MRSA	Ready to use	30 sec
Vancomycin-Resistant Enterococci (VRE)	Ready to use	30 sec
Candida albicans	Ready to use	30 sec
Mycobacterium terrae	Ready to use	30 sec
Mycobacterium avium	Ready to use	30 sec
All enveloped viruses (incl. HIV, HBV, HCV, Vaccinia)	Ready to use	15 sec
Noro virus	Ready to use	15 sec
Adeno virus	Ready to use	30 sec

BACTICID AF is tested according to the following European standards: EN 13624, EN 13697, EN 13727, EN 14348, EN 14476, EN 14561, EN 14562, EN 14563.

#### **INGREDIENTS:**

100 g of product contains active ingredients: ethanol 57 g, isopropanol 6 g.

### PRECAUTIONS:

Highly flammable liquid and vapour.

Keep out of reach of children. Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.

Keep container tightly closed. Wear protective gloves

Store in a well-ventilated place. Keep cool

Dispose of contents and / or containers in accordance with regulations on hazardous waste or packaging and packaging waste respectively

## **PACKAGING:**

0,5L	spray bottle
1 L	spray bottle
5 L	canister

AS Chemi-Pharm, Pöllu 132, Tallinn, 109117, Estonia, www.chemi-pharm.com