


Sudan Black B kit

Four-reagent kit for staining neutrophil granules in hematological smears (blood or bone marrow films). Used as one of the methods for detecting myelocytic and myelomonocytic leukemia.

for 350-400 tests SBBH-K-100


•



Sudan Black B Lipid kit

Four-reagent kit for specific lipid staining in cytochemistry. Sudan Black dye stains a few types of lipids, including neutral fats, phospholipids and sterols. Contains double amount of Sudan Black B, solution.


for 100 tests SBB-100T 5x100 mL SBB-K-100



TB-Stain Histo kit

Three-reagent kit for staining acid-fast bacteria (pathogenic mycobacteria) in histology sections, sputum, smears and culture smears according to Ziehl-Neelsen. Heating of the carbol-fuchsin solution is avoided in this protocol hence the release of hazardous phenolic vapors is omitted.


for 100 tests TBHK-100T



UriGnost S kit

Kit for sampling, staining, and microscopic analysis of urine sediment. Contains modified reagent according to Sternheimer (reagent manufactured according to the European Confederation of Laboratory Medicine (ECLM) and all the necessary equipment for sampling, concentrating, staining, and counting cells and kidney cylinders, as well as urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 μL pipette tips and urine plates).


for 500 tests USK-500 * for more information, see pages 57 and 75



UriGnost SM kit

Kit for sampling, staining and microscopic analysis of urine sediments. Contains modified reagent according to Sternheimer-Malbin and equipment required for sampling, concentration, staining, counting of kidney cells and kidney cylinders and urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 µL pipette tips and urine plates).

for 500 tests USMK-500 * for more information, see pages 57 and 76



Van Gieson Trichrome kit

Three-reagent kit for staining collagen fibers, muscle tissue, keratinized epithelium, cytoplasm, glial fibers and erythrocytes. Used for differentiation between collagen and smooth fibers in tumors and various other diseases.