OXOID QUALITY ASSURANCE PRODUCT SPECIFICATION

BROAD SPECTRUM BETA-LACTAMASE MIXTURE SR0113E

BROAD SPECTRUM BETA-LACTAMASE MIXTURE

SR0113E

Formula

Per vial	
Lyophilised β-Lactamases	
β-Lactamase I minimum	500 IU
β-Lactamase II minimum	50 IU

1 unit of enzyme activity will hydrolyse 1.0μ mole of substrate per minute at pH 7.0 and at 25° C. Note that 1 IU of activity = 600 Levy units of activity.

Description

The enzyme mixture is presented as a freeze-dried powder containing buffer and zinc salts.

Directions

Aseptically add 5ml of sterile distilled water into a vial of enzyme mixture and mix gently. Add 1ml of this solution aseptically to the blood culture bottle and mix well, preferably before or immediately after inoculation with the blood sample (5-10ml).

Physical Characteristics

Sterility - passes test

Stability of Reagents

Reconstituted solution of the enzyme can be stored at 2-8oC for up to four weeks. It can also be stored at -20oC, however, it is not advisable to store the solution for long periods because of the possibility of contamination. Repeated freezing and thawing should be avoided.

OXOID QUALITY ASSURANCE PRODUCT SPECIFICATION

BROAD SPECTRUM BETA-LACTAMASE MIXTURE SR0113E

Revision History

Section / Step	Description of Change	Reason for Change	Reference
Stability of	Update of the statement for	Updated togive clarification	MOC-2022-
Reagents	stability of reagents.		0362