



Technical Data

T.S.C. Supplement (Perfringens T.S.C. Supplement)

FD014

An antibiotic supplement, recommended for the selective isolation of Clostridium perfringens.

Composition

Per vial sufficient for 500 ml medium

*Ingredients

D-Cycloserine

Concentration

200mg

Directions:

Rehydrate the contents of 1 vial aseptically with 2 ml sterile distilled water. Mix well and aseptically add to 475 ml of sterile, molten, cooled (45-50°C) Perfringens Agar Base (T.S.C. / S.F.P., [M837](#)), Perfringens HiVeg™ Agar Base (T.S.C. / S.F.P., [MV837](#)), along with 25 ml of Egg Yolk Emulsion [FD045](#) or 500 ml of Perfringens Agar Base [M837I](#) / Tryptose Cycloserine Dextrose Agar Base [M1233](#) / Tryptose Cycloserine Dextrose HiVeg™ Agar Base [MV1233](#) / Tryptose Cycloserine Azide Agar Base [M1279](#). Tryptone Yeast Sodium sulphite Agar Base [M2046I](#)

Mix well and pour into sterile petri plates.

Storage and Shelf Life

Store at 2-8°C. Use before expiry date on the label.

Revision : 02 / 2017

* Not For Medicinal Use

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.