

Technical Data

Fraser Supplement

FD141

Recommended by ISO Committee for the selective isolation and enumeration of Listeria monocytogenes from food, animal feeds etc.

Composition

Per vial sufficient for 500 ml medium

IngredientsConcentrationFerric ammonium citrate250mg

Directions:

Rehydrate the contents of 1 vial aseptically with 1-2 ml of sterile distilled water. Mix well. For primary enrichment aseptically add 2 vials of FD141 to 1000 ml, sterile, cooled (45-50°C) Fraser Broth Base M1327 / Fraser HiVegTM Broth Base MV1327 along with rehydrated contents of 1 vial of Fraser Selective Supplement FD125I or add 1 vial of FD141 to 500 ml sterile Fraser Broth Base MV1327 along with rehydrated contents of 1 vial of Fraser Selective Supplement FD125I for secondary enrichment. Mix well before dispensing as desired.

Storage and Shelf Life

Store between 10-30°C. Use before the expiry date on the label.

Revision: 1/2012

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia[™] publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia[™] Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.