



DR. BRILL + DR. STEINMANN
INSTITUTE FOR HYGIENE AND MICROBIOLOGY



20/02/2018

Test report L17/0629aMV.2

Evaluation of the effectiveness of **Bactid AF**

Test virus: modified vaccinia virus Ankara (MVA)

Method: based on EN 14476:2013+A1:2015 (dirty conditions)

quantitative suspension test for the evaluation
of virucidal activity of chemical disinfectants and
antiseptics used in human medicine

Sponsor:
Chemi-Pharm AS
Pollu 132
EST – TALLINN 10917

Norderoog 2, DE - 28259 Bremen
Tel.: +49 40-557631-0, Fax: +49 40-557631-11
info@brillhygiene.com, <http://www.brillhygiene.com>

1. Identification of test laboratory

Dr. Brill + Partner GmbH Institute for Hygiene and Microbiology, Norderoog 2, DE - 28259 Bremen

2. Identification of sample

Manufacturer	Chemi-Pharm AS
Name of product	Bactacid AF
Confirmation no.	203850
Product diluent recommended by the manufacturer	-
Batch number	197101017
Application	surface disinfection
Production date	10/10/2017
Expiry date	10/10/2020
Active compound (s) (100 g)	57 g ethanol 6 g IPA
Appearance, odour	clear, colorless liquid product specific
pH-values	undiluted: 7.49 (20 °C)
Storage conditions	room temperature in the dark (area with restricted access)
Date of arrival in the laboratory	13/10/2017

3. Materials

3.1 Culture medium and reagents

- Eagle's Minimum Essential Medium with Hank's BSS (MEM, Biozym Scientific GmbH, catalogue no. 880144)
- fetal calf serum (Biochrom AG, article no. S 0115)
- 1.4 % formaldehyde solution (dilution of Roti®-Histofix 4 %, Carl Roth GmbH)
- Aqua bidest. (SG ultrapure water system, type Ultra Clear; serial no. 86996-1)
- PBS (Invitrogen, article no. 18912-014)

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

- BSA (Sigma-Aldrich-Chemie GmbH, article no. CA-2153)
- sheep erythrocytes (Fiebig Nährstofftechnik).

3.2 Virus and cells

The modified vaccinia virus Ankara (MVA) originated from Dr. Manteufel, Institut für Tierhygiene und Öffentliches Veterinärwesen, DE - 04103 Leipzig. Before inactivation assays, virus had been passaged three times in *BHK 21-cells* (Baby Hamster Kidney).

BHK 21-cells (passage 108) originated from the Friedrich-Löffler-Institut, Bundesforschungsinstitut für Tiergesundheit (formerly Bundesforschungsanstalt für Viruskrankheiten der Tiere, Isle of Riems).

The cells were inspected regularly for morphological alterations and for contamination by mycoplasmas. No morphological alterations of cells and no contamination by mycoplasmas could be detected.

3.3 Apparatus, glassware and small items of equipment

- CO₂ incubator, Nunc GmbH & Co. KG, model QWJ 350
- Agitator (Vortex Genie Mixer, type G 560E)
- pH measurement 315i (WTW, article no. 2A10-100)
- Centrifuge (Sigma-Aldrich-Chemie GmbH, type 113)
- Microscope (Olympus, type CK 30)
- Centrifuge 5804 R (Eppendorf AG)
- Water bath (JULABO, Julabo U 3)
- Adjustable and fixed-volume pipettes (Eppendorf AG)
- Polystyrene 96-well microtitre plate (Nunc GmbH & Co. KG, Wiesbaden)
- Cell culture flask (Nunc GmbH & Co. KG, Wiesbaden)
- Sealed test tubes (Sarstedt AG & Co., Nümbrecht).

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE - 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

4. Experimental conditions

Test temperature	20 °C ± 1.0 °C
Concentration of test product	undiluted (80.0 %) and as 50.0 % and 10.0 % (demonstration of non-active range) solutions
Appearance of product dilutions	no precipitation
Contact times	15, 20, 30 and 60 seconds and 30 minutes
Interfering substance	3.0 g/l bovine serum albumin + 3.0 g/l erythrocytes (dirty conditions, EN 14476)
Procedure to stop action of disinfectant	immediate dilution
Diluent	water
Stability of product in the mix with virus and interfering substance (80.0 % solution)	minor clouding, no precipitation
Virus strain	modified vaccinia virus Ankara (MVA) (ATCC VR-1508)
Date of testing	19/12/2017 – 20/02/2018
End of testing	20/02/2018

5. Methods

5.1 Preparation of test virus suspension

For preparation of test virus suspension, *BHK 21-cells* were cultivated with MEM and 10 % or 2 % fetal calf serum. Cells were infected with a multiplicity of infection of 0.1. After cells showed a cytopathic effect, they were subjected to a freeze/thaw procedure followed by a low speed centrifugation in order to sediment cell debris. After aliquotation, test virus suspension was stored at – 80 °C.

5.2 Preparation of disinfectant (dilutions)

The test product was tested undiluted. Due to the addition of interfering substance and test virus suspension an 80.0 % solution resulted.

Furthermore, the product was evaluated as 50.0 % and 10.0 % solutions (demonstrating of non-active range). These solutions were prepared with water immediately before the inactivation tests.

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

5.3 Infectivity assay

Infectivity was determined as endpoint titration according to EN 5.5 transferring 0.1 ml of each dilution into eight wells of a microtitre plate to 0.1 ml of freshly trypsinised *BHK 21-cells* ($10\text{--}15 \times 10^3$ cells per well), beginning with the highest dilution. Microtitre plates were incubated at 37 °C in a 5 % CO₂-atmosphere. The cytopathic effect was read by using an inverted microscope after six days. Calculation of the infective dose TCID₅₀/ml was calculated with the method of Spearman (2) and Kärber (3) with the following formula:

$$-\log_{10}\text{TCID}_{50} = X_0 - 0.5 + \sum r/n$$

meaning

X_0 = log₁₀ of the lowest dilution with 100 % positive reaction

r = number of pos. determinations of lowest dilution step with 100 % positive and all higher positive dilution steps

n = number of determinations for each dilution step.

5.4 Calculation and verification of virucidal activity

The virucidal activity of the test disinfectant was evaluated by calculating the decrease in titre in comparison with the control titration without disinfectant. The difference is given as reduction factor (RF).

According to the EN 14476, a disinfectant or a disinfectant solution at a particular concentration is having virus-inactivating efficacy if the titre is reduced at least by 4 log₁₀ steps within the recommended exposure period. This corresponds to an inactivation of ≥ 99.99 %.

5.5 Inactivation assay

Determination of virucidal activity has been carried out based on EN 5.5. The test product was examined undiluted (80.0 %) and as 50.0 % and 10.0 % (demonstration of non-active range) solutions in water at 20 °C based on EN 14476. 15, 20, 30 and 60 seconds and 30 minutes were chosen as contact times.

Immediately at the end of a chosen contact time, activity of the disinfectant was stopped by dilution to 10⁻⁸.

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

Titration of the virus control was performed at the beginning of the test and after the longest exposure time (EN 5.5.7). One part by volume of test virus suspension was mixed with one part interfering substance and eight parts by volume of WSH or Aqua bidest. (RTU products).

Furthermore, a cell control (only addition of medium) was incorporated.

Inactivation tests were carried out in sealed test tubes in a water bath at $20\text{ °C} \pm 1.0\text{ °C}$. Aliquots were retained after appropriate exposure times and residual infectivity was determined.

5.6 Determination of cytotoxicity

Determination of cytotoxicity was performed according to EN 5.5.4.1.

5.7 Cell sensitivity to virus

For the control of cell sensitivity to virus two parts by volume of water were mixed with eight parts by volume of the lowest apparently non-cytotoxic dilution of the product. These mixtures or PBS as control were added to a volume of double concentrated cell suspension. After 1 h at 37 °C the cells were centrifuged and re-suspended in cell culture medium (EN 5.5.4.2b).

Finally, a comparative titration of the test virus suspension was performed on the pre-treated (disinfectant) and non-pre-treated (PBS) cells as described above.

5.8 Control of efficacy for suppression of disinfectant's activity

Furthermore, a control of efficiency for suppression of disinfectant's activity was included (EN 5.5.5).

5.9 Reference virus inactivation test

As reference for test validation a 0.7 % formaldehyde solution according to EN 5.5.6 was included. 5, 15, 30 and 60 minutes were chosen as contact times. In addition, cytotoxicity of formaldehyde test solution was determined according to EN 5.5.6.2 with dilutions up to 10^{-5} .

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

6. Verification of the methodology

The following criteria as mentioned in EN 5.7 were fulfilled:

- a) The titre of the test virus suspension allowed the determination of a $\geq 4 \log_{10}$ reduction (maximal virus reduction $\geq 5.00 \pm 0.33$)
- b) The test product (80.0 %) showed no cytotoxicity in the 1:10 dilutions thus allowing the detection of a $4 \log_{10}$ reduction of virus titre.
- c) The comparative titration on pre-treated (disinfectant) and non-pre-treated (PBS) *BHK 21-cells* showed no significant difference ($< 1 \log_{10}$; EN 5.7) of virus titre: 6.50 ± 0.00 (PBS) versus 6.25 ± 0.33 (1:10 dilution of disinfectant as 80.0 % solution) $\log_{10}$ TCID₅₀/ml.
- d) The control of efficacy for suppression of disinfectant's activity (80.0 %) showed no decrease ($\leq 0.5 \log_{10}$; EN 5.5.5.1) in virus titre (6.63 ± 0.41 versus $6.50 \pm 0.46 \log_{10}$ TCID₅₀/ml).
- e) One concentration demonstrated a $4 \log_{10}$ reduction and (at least) one concentration demonstrated a $\log_{10}$ reduction of less than 4.

Since all criteria according EN 5.7 were fulfilled, examination with MVA based on EN 14476 is valid.

7. Results

Results of examination are shown in tables 1 to 8. Tables 1 to 7 demonstrate the raw data, whereas table 8 (a+b) gives a summary of results.

The undiluted test product as 80.0 % solution was able to inactivate MVA after 15 seconds of exposure time under dirty conditions (tables 1 and 2). No residual virus was found at this time point. The reduction factors were $\geq 4.88 \pm 0.29$ and $\geq 5.00 \pm 0.33$ (mean value $\geq 4.94 \pm 0.22$). This corresponded to an inactivation of ≥ 99.99 %.

Tested as 50.0 % solution, the product was not active within 30 seconds of exposure time (table 3). The reduction factor was 1.13 ± 0.53 .

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

Tested as 10.0 % solution, the product was not active within 30 minutes of exposure time (table 4).

8. Conclusion

The surface disinfectant Bactacid AF tested undiluted demonstrated activity against MVA after an exposure time of 15 seconds under dirty conditions.

Therefore, the surface disinfectant Bactacid AF can be declared as active against MVA as follows:

undiluted	15 seconds	dirty conditions
-----------	------------	------------------

Bremen, 20/02/2018



- Dr. Britta Becker -
Head of Laboratory



- Dr. Dajana Paulmann -
Scientific Project Manager



*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

9. Quality control

The Quality Assurance of the results was maintained by performing the determination of the virus-inactivating properties of the disinfectant in accordance with Good Laboratory Practice regulations:

- 1) Chemicals Act of Germany, Appendix 1, dating of 01.08 1994 (BGBl. I, 1994, page 1703). Appendix revised at 14. 05. 1997 (BGBl. I, 1997, page 1060).
- 2) OECD Principles of Good Laboratory Practice (revised 1997); OECD Environmental Health and Safety Publications; Series on Principles of Good Laboratory Practice and Compliance Monitoring – Number 1. Environment Directorate, Organization for Economic Co-operation and Development, Paris 1998.

The plausibility of the results was additionally confirmed by controls incorporated in the inactivation assays.

10. Records to be maintained

All testing data, protocol, protocol modifications, the final report, and correspondence between Dr. Brill + Partner GmbH and the sponsor will be stored in the archives at Dr. Brill + Partner GmbH.

The use of the Dr. Brill + Partner GmbH name, logo or any other representation of Dr. Brill + Partner GmbH, other than distribution of this report in it's entirely, without the written approval of Dr. Brill + Partner GmbH is prohibited. In addition, Dr. Brill + Partner GmbH may not be referred to in any form of promotional materials, press releases, advertising or similar materials (whether by print, broadcast, communication or electronic means) without the express permission of Dr. Brill + Partner GmbH.

The test results in this test report relate only to the items examined.

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

11. Literature

1. EN 14476:2013+A1:2015: Chemical disinfectants and antiseptics – Quantitative suspension test for the evaluation of virucidal activity of chemicals disinfectants and antiseptics in human medicine test - Test method and requirements (phase 2, step 1)
2. Spearman, C.: The method of 'right or wrong cases' (constant stimuli) without Gauss's formulae.
Brit J Psychol; 2 1908, 227-242
3. Kärber, G.: Beitrag zur kollektiven Behandlung pharmakologischer Reihenversuche.
Arch Exp Path Pharmac; 162, 1931, 480-487

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018

Appendix:

Legend to the Tables

Table 1:	Raw data for Bactacid AF (80.0 %) tested against MVA (1 st assay)
Table 2:	Raw data for Bactacid AF (80.0 %) tested against MVA (2 nd assay)
Table 3:	Raw data for Bactacid AF (50.0 %) tested against MVA
Table 4:	Raw data for Bactacid AF (10.0 %) tested against MVA
Table 5:	Raw data for formaldehyde solution (0.7 %) tested against MVA
Table 6:	Raw data for control of efficacy for suppression of disinfectant's activity (80.0 %)
Table 7:	Raw data (MVA) for cell sensitivity (80.0 %)
Table 8 (a+b):	Summary of results with Bactacid AF and MVA

Legend to the Figures

Figure 1:	Virus-inactivating properties of Bactacid AF (80.0 %)
Figure 2:	Virus-inactivating properties of formaldehyde (0.7 %)

*Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderoog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018



Table 1: Raw data for BactiCid AF (80.0 %) tested against MVA at 20 °C (quantal test; 8 wells) (#5390) (1st assay)

Product	Concentration	Interfering substance	Contact time	Dilutions (log ₁₀)								
				1	2	3	4	5	6	7	8	9
test product	80.0 %	dirty conditions	15 s	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.
			20 s	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.
			30 s	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.
			60 s	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.
test product cytotoxicity	80.0 %	dirty conditions	n.a.	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d. 0000	n.d.	n.d.	n.d.
virus control	n.a.	dirty conditions	0 min	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			60 min	4444 4444	4444 4444	4444 4444	4433 3333	3202 2202	0000 2000	0000 0000	0000 0000	0000 0000

n.a. = not applicable
n.d. = not done

0 = no virus present; t = cytotoxic

1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 2: Raw data for BactiCid AF (80.0 %) tested against MVA at 20 °C (quantal test; 8 wells) (#5404) (2nd assay)

Product	Concentration	Interfering substance	Contact time	Dilutions (log ₁₀)								
				1	2	3	4	5	6	7	8	9
test product	80.0 %	dirty conditions	15 s	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.
			20 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			30 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			60 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
test product cytotoxicity	80.0 %	dirty conditions	n.a.	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.	
virus control	n.a.	dirty conditions	0 min	4444 4444	4444 4444	4444 4444	4444 4444	3223 3443	0000 2000	0000 0000	0000 0000	
			60 min	4444 4444	4444 4444	4444 4444	4444 4444	4420 0242	0000 2020	0000 0000	0000 0000	

n.a. = not applicable

0 = no virus present; t = cytotoxic

n.d. = not done

1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 3: Raw data for Bactcid AF (50.0 %) tested against MVA at 20 °C (quantal test; 8 wells) (#5390)

Product	Concentration	Interfering substance	Contact time	Dilutions (log ₁₀)									
				1	2	3	4	5	6	7	8	9	
test product	50.0 %	dirty conditions	15 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			20 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			30 s	4444 4444	4444 4444	3333 3323	2200 3222	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.
			60 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
test product cytotoxicity	50.0 %	dirty conditions	n.a.	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.	n.d.	n.d.
virus control	n.a.	dirty conditions	0 min	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			60 min	4444 4444	4444 4444	4444 4444	4433 3333	3202 2202	0000 2000	0000 0000	0000 0000	0000 0000	0000 0000

n.a. = not applicable
n.d. = not done

0 = no virus present; t = cytotoxic
1 to 4 = virus present (degree of CPE)

1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 4: Raw data for BactiCid AF (10.0 %) tested against MVA at 20 °C (quantal test; 8 wells) (#5390)

Product	Concentration	Interfering substance	Contact time	Dilutions (log ₁₀)								
				1	2	3	4	5	6	7	8	9
test product	10.0 %	dirty conditions	15 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			20 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			30 s	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			30 min	4444 4444	4444 4444	4444 4444	3433 3344	0202 1001	0000 0000	0000 0000	0000 0000	n.d.
test product cytotoxicity	10.0 %	dirty conditions	n.a.	0000 0000	0000 0000	0000 0000	0000 0000	n.d.	n.d.	n.d.	n.d.	
virus control	n.a.	dirty conditions	0 min	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	
			60 min	4444 4444	4444 4444	4444 4444	4433 3333	3202 2202	0000 2000	0000 0000	0000 0000	0000 0000

n.a. = not applicable
n.d. = not done

0 = no virus present; t = cytotoxic

1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 5: Raw data for formaldehyde solution (0.7 %) tested against MVA at 20 °C (quantal test; 8 wells) (#5404)

Product	Concentration	Interfering substance	Contact time (min)	Dilutions (log ₁₀)								
				1	2	3	4	5	6	7	8	9
formaldehyde	0.7 % (m/V)	PBS	5	tttt tttt	tttt tttt	tttt tttt	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d. n.d.
			15	tttt tttt	tttt tttt	tttt tttt	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d. n.d.
			30	tttt tttt	tttt tttt	tttt tttt	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d. n.d.
			60	tttt tttt	tttt tttt	tttt tttt	0000 0000	0000 0000	0000 0000	0000 0000	0000 0000	n.d. n.d.
formaldehyde cytotoxicity	0.7 % (m/V)	PBS	n.a.	tttt tttt	tttt tttt	tttt tttt	0000 0000	0000 0000	n.d.	n.d.	n.d.	n.d.
virus control	n.a.	PBS	0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
			60	4444 4444	4444 4444	4444 4444	4444 4444	3233 2333	2000 0000	0000 0000	0000 0000	0000 0000

n.a. = not applicable
n.d. = not done

0 = no virus present; t = cytotoxic
1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 6: Raw data for control of efficacy for suppression of disinfectant's activity (80.0 %) (#5404)

Product	Interfering substance	dilutions (log ₁₀)								
		1	2	3	4	5	6	7	8	9
test product	dirty conditions	4444 4444	4444 4444	4444 4444	4444 4444	3434 4044	0020 0002	0000 0000	0000 0000	n.d.
corresponding virus control	dirty conditions	4444 4444	4444 4444	4444 4444	4444 4444	4420 0242	0000 2020	0000 0000	0000 0000	0000 0000

n.a. = not applicable

0 = no virus present; t = cytotoxic

n.d. = not done

1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 7: Raw data (MVA) for cell sensitivity (80.0 % solution) (#5404)

Product	Dilution	Dilutions (log ₁₀)								
		1	2	3	4	5	6	7	8	9
PBS	-	4444	4444	4444	4444	2223	0000	0000	0000	n.d.
		4444	4444	4444	4444	3234	0000	0000	0000	
test product	1:10	4444	4444	4444	4444	2430	0000	0000	0000	n.d.
		4444	4444	4444	4444	3203	0000	0000	0000	

n.a. = not applicable

0 = no virus present; t = cytotoxic

n.d. = not done

1 to 4 = virus present (degree of CPE in 8 cell culture units) (wells of microtitre plates)



Table 8a: Summary of results with Bactioid AF and MVA

Product*	Con- centration	Interfering substance	Level of cytotoxicity	log ₁₀ TCID ₅₀ /ml after					> 4 log ₁₀ reduction after ...
				15 s	20 s	30 s	60 s	30 min	
test product (1)	80.0 %	dirty conditions	1.50	≤ 1.50±0.00	≤ 1.50±0.00	≤ 1.50±0.00	≤ 1.50±0.00	n.d.	15 s (RF ≥ 4.88±0.29)
test product (2)	80.0 %	dirty conditions	1.50	≤ 1.50±0.00	n.d.	n.d.	n.d.	n.d.	15 s (RF ≥ 5.00±0.33)
test product (2)	50.0 %	dirty conditions	1.50	n.d.	n.d.	5.25±0.33	n.d.	n.d.	> 30 s (RF = 1.13±0.33)
test product (2)	10.0 %	dirty conditions	1.50	n.d.	n.d.	n.d.	n.d.	6.00±0.38	> 30 min (RF = 0.38±0.29)

n.a. = not applicable n.d. = not done

*The number in brackets gives the number of the corresponding virus control (see table 8b)




Table 8b: Summary of results with Bactacid AF and MVA

Product	Con- centration	Interfering substance	Level of cytotoxicity	log ₁₀ TCID ₅₀ /ml aftermin					> 4 log ₁₀ reduction after ... min
				0	5	15	30	60	
formaldehyde	0.7 % (w/v)	PBS	4.50	n.d.	≤ 4.50±0.00	≤ 4.50±0.00	≤ 4.50±0.00	≤ 4.50±0.00	≥ 5 (RF ≥ 2.13±0.25)
virus control	n.a.	PBS	n.a.	n.d.	n.d.	n.d.	n.d.	6.63±0.25	n.a.
virus control (1)	n.a.	dirty conditions	n.a.	n.d.	n.d.	n.d.	n.d.	6.38±0.41	n.a.
virus control (2) (+ suppression)	n.a.	dirty conditions	n.a.	6.63±0.25	n.d.	n.d.	n.d.	6.50±0.46	n.a.
suppression control	80.0 %	dirty conditions	1.50	n.d.	n.d.	n.d.	6.38±0.41	n.d.	n.a.
sens. PBS	n.a.	n.a.	n.a.	n.d.	n.d.	n.d.	n.d.	6.50±0.00	n.a.
sens. product	80.0 % → 1:10	n.a.	n.a.	n.d.	n.d.	n.d.	n.d.	6.25±0.33	n.a.

n.a. = not applicable n.d. = not done sens. = sensitivity




Figure 1: Virus-inactivating properties of Bactacid AF (80.0 %)



* Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Norderpoog 2, DE – 28259 Bremen, Germany, Telephone +49 40 557631-0, Telefax +49 40 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018



Figure 2: Virus-inactivating properties of formaldehyde (0.7 %)



* Test procedure accredited according to DIN EN ISO/IEC 17025. Test report issued by Dr. Brill + Partner GmbH, Nordroog 2, DE – 28259 Bremen, Germany, Telephone +49. 40. 557631-0, Telefax +49. 40. 557631-11, www.brillhygiene.com. No copying or transmission, in whole or in part, of this test report without the explicit prior written permission. The test results exclusively apply to the tested samples. Information on measurement uncertainty on request. © Dr. Brill + Partner GmbH 2018