



Technical Data

SS Agar (Salmonella Shigella Agar)

M108

Intended Use:

Recommended for the isolation of *Salmonella* and some *Shigella* species from pathological specimens, suspected foodstuffs etc.

Composition**

Ingredients	Gms / Litre
Peptone	5.000
HM peptone B #	5.000
Lactose	10.000
Bile salts mixture	8.500
Sodium citrate	10.000
Sodium thiosulphate	8.500
Ferric citrate	1.000
Brilliant green	0.00033
Neutral red	0.025
Agar	15.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

- Equivalent to Beef extract

Directions

Suspend 63.02 grams in 1000 ml purified /distilled water. Boil with frequent agitation to dissolve the medium completely. DO NOT AUTOCLAVE OR OVERHEAT. Overheating may destroy selectivity of the medium. Cool to about 50°C. Mix and pour into sterile Petri plates.

Principle And Interpretation

SS Agar medium is recommended as differential and selective medium for the isolation of *Salmonella* and *Shigella* species from pathological specimens (4) and suspected foodstuffs (1,6, 8, 9) and for microbial limit test (7). SS Agar is a moderately selective medium in which gram-positive bacteria are inhibited by bile salts, brilliant green and sodium citrate.

Peptone, HM peptone B provides nitrogen and carbon source, long chain amino acids, vitamins and essential growth nutrients. Lactose is the fermentable carbohydrate. Brilliant green, bile salts and thiosulphate selectively inhibit gram-positive and coliform organisms. Sodium thiosulphate is reduced by certain species of enteric organisms to sulphite and H₂S gas and this reductive enzyme process is attributed by thiosulphate reductase. Production of H₂S gas is detected as an insoluble black precipitate of ferrous sulphide, formed upon reaction of H₂S with ferric ions or ferric citrate, indicated in the center of the colonies.

The high selectivity of Salmonella Shigella Agar allows the use of large inocula directly from faeces, rectal swabs or other materials suspected of containing pathogenic enteric bacilli. On fermentation of lactose by few lactose-fermenting normal intestinal flora, acid is produced which is indicated by change of colour from yellow to red by the pH indicator-neutral red. Thus these organisms grow as red pigmented colonies. Lactose non-fermenting organisms grow as translucent colourless colonies with or without black centers. Growth of *Salmonella* species appears as colourless colonies with black centers resulting from H₂S production. *Shigella* species also grow as colourless colonies which do not produce H₂S.

Type of specimen

Clinical: faeces, blood, rectal swabs; Suspected food stuffs.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (2,3,4).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,6,8,9).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. The medium is highly selective and may be toxic to certain *Salmonella* or *Shigella* species. Hence it is recommended to use to inoculate plates of less inhibitory media parallel to SS Agar, such as Hektoen Enteric Agar (M467) or Deoxycholate Citrate Agar (M065) for easier isolation of *Shigella* species (6).

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Reddish orange coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 6.3% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
# <i>Klebsiella aerogenes</i> ATCC 13048 (00175*)	50-100	fair	20-30%	cream pink
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	fair	20-30%	pink with bile precipitate
<i>Salmonella Choleraesuis</i> ATCC 12011	50-100	good-luxuriant	≥50%	colourless with black centre
<i>Salmonella Typhi</i> ATCC 6539	50-100	good-luxuriant	≥50%	colourless with black centre
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	50-100	none-poor	≤10%	colourless
<i>Proteus mirabilis</i> ATCC 25933	50-100	fair-good	30-40%	colourless, may have black centre
<i>Shigella flexneri</i> ATCC 12022 (00126*)	50-100	good	40-50%	colourless
<i>Salmonella Typhimurium</i> ATCC 14028 (00031*)	50-100	good-luxuriant	≥50%	colourless with black centre
<i>Salmonella Enteritidis</i> ATCC 13076 (00030*)	50-100	good-luxuriant	≥50%	colourless with black centre

Key : *Corresponding WDCM numbers.

Formerly known as *Enterobacter aerogenes*

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

1. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
4. Lennette and others (Eds.), 1985, Manual of Clinical Microbiology, 4th ed., ASM, Washington, D.C.
5. MacFaddin J., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
6. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
7. The United States Pharmacopoeia, 2006, USP29/NF24, The United States Pharmacopoeial Convention. Rockville, MD.
8. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
9. Williams S., (Ed.), 2005, Official Methods of Analysis of the Association of Official Analytical Chemists, 19th Ed., AOAC, Washington, D.C.


Revision : 03/
2018



In vitro diagnostic medical
device



CE Marking



Storage temperature



Do not use if package is
damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India



CE Partner 4U, Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.