

Formulae

Difco™ MYP Agar

Approximate Formula* Per 900 mL

Beef Extract.....	1.0	g
Peptone.....	10.0	g
D-Mannitol.....	10.0	g
Sodium Chloride.....	10.0	g
Phenol Red.....	25.0	mg
Agar.....	15.0	g

Difco™ Antimicrobial Vial P

Approximately 30,000 units polymyxin B per vial.

*Adjusted and/or supplemented as required to meet performance criteria.

Directions for Preparation from Dehydrated Product

Difco™ MYP Agar

1. Suspend 46 g of the powder in 900 mL of purified water. Mix thoroughly.
2. Heat with frequent agitation and boil for 1 minute to completely dissolve the powder.
3. Dispense 225 mL into 500 mL flasks.
4. Autoclave at 121°C for 15 minutes. Cool to 45-50°C.
5. Aseptically add 12.5 mL Egg Yolk Enrichment 50% and 4.1 mL Antimicrobial Vial P rehydrated with 5 mL sterile water (25,000 units of polymyxin B). Mix thoroughly.
6. Test samples of the finished product for performance using stable, typical control cultures.

Difco™ Antimicrobial Vial P (Polymyxin B)

1. To rehydrate, aseptically add 5 mL sterile purified water (to achieve the desired concentration for MYP Agar).
2. Rotate in an end-over-end motion to dissolve the contents completely.

Procedure

Consult appropriate references.^{4,6}

Expected Results

Consult appropriate references.^{4,6}

References

1. Mosse, Koopman and Jongerius. 1967. *Appl. Microbiol.* 15:650.
2. Donovan. 1958. *J. Appl. Bacteriol.* 21:100.
3. Coliner. 1948. *J. Bacteriol.* 55:777.
4. U.S. Food and Drug Administration. 2001. *Bacteriological analytical manual*, online. AOAC International, Gaithersburg, Md.
5. Bennett and Belay. 2001. *In* Downes and Ito (ed.), *Compendium of methods for the microbiological examination of foods*, 4th ed. American Public Health Association, Washington, D.C.
6. Horwitz (ed.). 2007. *Official methods of analysis of AOAC International*, 18th ed., online. AOAC International, Gaithersburg, Md.

Availability

Difco™ MYP Agar

AOAC BAM COMPF ISO USDA

Cat. No. 281010 Dehydrated – 500 g

Europe

Cat. No. 257004 Prepared Plates – Pkg. of 20*

Japan

Cat. No. 251264 Prepared Plates – Pkg. of 20*

Difco™ Antimicrobial Vial P

AOAC BAM COMPF ISO USDA

Cat. No. 232681 Vial – 6 × 10 mL*

Difco™ Egg Yolk Enrichment 50%

AOAC BAM COMPF ISO USDA

Cat. No. 233471 Tube – 12 × 10 mL*

233472 Bottle – 6 × 100 mL*

*Store at 2-8°C.

MacConkey Agars

MacConkey Agar • MacConkey Agar Base

MacConkey Agar without Crystal Violet

MacConkey Agar without Crystal Violet or Salt

MacConkey Agar without Salt

Intended Use

MacConkey agars are slightly selective and differential plating media mainly used for the detection and isolation of gram-negative organisms from clinical,¹⁻³ dairy,⁴ food,⁵⁻⁷ water,⁸ pharmaceutical,⁹⁻¹¹ cosmetic,^{6,7} and other industrial sources.

MacConkey Agar is used for isolating and differentiating lactose-fermenting from lactose-nonfermenting gram-negative enteric bacilli.

MacConkey Agar Base is used with added carbohydrate in differentiating coliforms based on fermentation reactions.

MacConkey Agar without Crystal Violet is used for isolating and differentiating enteric microorganisms while permitting growth

of staphylococci and enterococci. The medium can be used also to separate *Mycobacterium fortuitum* and *M. chelonae* from other rapidly growing mycobacteria.

MacConkey Agar without Crystal Violet or Salt and MacConkey Agar without Salt are used for isolating and differentiating gram-negative bacilli while suppressing the swarming of most *Proteus* species.

MacConkey Agar meets *United States Pharmacopeia (USP)*, *European Pharmacopoeia (EP)* and *Japanese Pharmacopoeia (JP)*⁹⁻¹¹ performance specifications, where applicable.

User Quality Control

NOTE: Differences in the Identity Specifications and Cultural Response testing for media offered as both **Difco™** and **BBL™** brands may reflect differences in the development and testing of media for industrial and clinical applications, per the referenced publications.

Identity Specifications

Difco™ MacConkey Agar

Dehydrated Appearance:	Pink to pinkish beige, free-flowing, homogeneous.
Solution:	5.0% solution, soluble in purified water upon boiling. Solution is reddish-purple, slightly opalescent.
Prepared Appearance:	Pinkish red, slightly opalescent.
Reaction of 5.0% Solution at 25°C:	pH 7.1 ± 0.2

Difco™ MacConkey Agar Base

Dehydrated Appearance:	Pinkish beige, free-flowing, homogeneous.
Solution:	4.0% solution, soluble in purified water upon boiling. Solution is red, very slightly to slightly opalescent.
Prepared Appearance:	Red, slightly opalescent.
Reaction of 4.0% Solution at 25°C:	pH 7.1 ± 0.2

Difco™ MacConkey Agar without Crystal Violet

Dehydrated Appearance:	Pinkish beige, free-flowing, homogeneous.
Solution:	5.2% solution, soluble in purified water upon boiling. Solution is reddish orange, clear to very slightly opalescent.
Prepared Appearance:	Reddish orange, clear to very slightly opalescent.
Reaction of 5.2% Solution at 25°C:	pH 7.4 ± 0.2

Difco™ MacConkey Agar without Salt

Dehydrated Appearance:	Beige to pinkish beige, free-flowing, homogeneous.
Solution:	4.7% solution, soluble in purified water upon boiling. Solution is reddish orange, slightly opalescent.
Prepared Appearance:	Reddish orange, slightly opalescent.
Reaction of 4.7% Solution at 25°C:	pH 7.4 ± 0.2

Cultural Response

Difco™ MacConkey Agar

Prepare the medium per label directions. Inoculate and incubate at 35 ± 2°C for 18-24 hours (incubate *E. coli* ATCC 25922 for 40-48 hours). For *E. coli* ATCC 8739, inoculate in duplicate and incubate one plate at 30-35°C for 18-24 hours and the other plate at 35-37°C for 18-72 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	29212	10 ³	Marked to complete inhibition	–	–
<i>Escherichia coli</i>	25922	30-300	Good	Pink to red	+
<i>Proteus mirabilis</i>	12453	30-300	Good	Colorless	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	30-300	Good	Colorless	–
<i>Escherichia coli</i>	8739	<100	Growth (18-24 hours at 30-35°C)	Pink to red	+
<i>Escherichia coli</i>	8739	<100	Growth (18-72 hours at 35-37°C)	Pink to red	+

Difco™ MacConkey Agar Base

Prepare the medium per label directions without and with 1% added lactose. Inoculate and incubate at 35 ± 2°C for 18-24 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR PLAIN	COLONY COLOR W/LACTOSE	BILE PPT.
<i>Enterococcus faecalis</i>	29212	10 ³	Marked to complete inhibition	–	–	–
<i>Escherichia coli</i>	25922	30-300	Good	Colorless	Pink to red (w/lactose)	+
<i>Proteus mirabilis</i>	12453	30-300	Good	Colorless	Colorless	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	30-300	Good	Colorless	Colorless	–

Difco™ MacConkey Agar without Crystal Violet

Prepare the medium per label directions. Inoculate and incubate at 35 ± 2°C for 18-48 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	29212	30-300	Good	Red	–
<i>Escherichia coli</i>	25922	30-300	Good	Pink to red	–
<i>Proteus mirabilis</i>	12453	30-300	Good	Colorless	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	30-300	Good	Colorless	–
<i>Staphylococcus aureus</i>	25923	30-300	Good	Pink to red	–

Difco™ MacConkey Agar without Salt

Prepare the medium per label directions. Inoculate and incubate at 35 ± 2°C for 18-48 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	33186	30-300	Good	Red	–
<i>Escherichia coli</i>	25922	30-300	Good	Pink to red	–
<i>Proteus mirabilis</i>	12453	30-300	Good	Colorless, no swarming	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	30-300	Good	Colorless	–
<i>Shigella flexneri</i>	12022	30-300	Good	Colorless	–

Continued

Identity Specifications**BBL™ MacConkey Agar**

Dehydrated Appearance: Fine, homogenous, may contain dark particles.

Solution: 5.0% solution, soluble in purified water upon boiling. Solution is medium to dark, rose to brown-rose with or without a trace orange tint; clear to slightly hazy.

Prepared Appearance: Medium to dark, rose to brown-rose with or without a trace orange tint; clear to slightly hazy.

Reaction of 5.0% Solution at 25°C: pH 7.1 ± 0.2

BBL™ MacConkey Agar (prepared)

Appearance: Medium-dark, rose-tan and trace hazy.

Reaction at 25°C: pH 7.1 ± 0.2

BBL™ MacConkey Agar without Crystal Violet

Dehydrated Appearance: Fine, homogeneous, free of extraneous material.

Solution: 5.2% solution, soluble in purified water upon boiling. Solution is medium, red-orange to red-rose, slightly hazy to hazy.

Prepared Appearance: Medium, red-orange to red-rose, slightly hazy to hazy.

Reaction of 5.2% Solution at 25°C: pH 7.4 ± 0.2

BBL™ MacConkey Agar without Crystal Violet or Salt

Dehydrated Appearance: Fine, homogeneous, free of extraneous material.

Solution: 4.37% solution, soluble in purified water upon boiling. Solution is medium, red-orange to red-rose, slightly hazy to hazy.

Prepared Appearance: Medium, red-orange to red-rose, slightly hazy to hazy.

Reaction of 4.37% Solution at 25°C: pH 7.4 ± 0.2

Cultural Response**BBL™ MacConkey Agar**

Prepare the medium per label directions. Inoculate and incubate at 35 ± 2°C for 48 hours. For *E. coli* ATCC 8739, inoculate in duplicate and incubate one plate at 30-35°C for 18-24 hours and the other plate at 35-37°C for 18-72 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	29212	10 ⁴ -10 ⁵	Partial to complete inhibition	–	–
<i>Escherichia coli</i>	25922	10 ³ -10 ⁴	Good	Red to rose-red	+
<i>Proteus mirabilis</i>	12453	10 ³ -10 ⁴	Good	Colorless	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	10 ³ -10 ⁴	Good	Colorless	–
<i>Shigella flexneri</i>	12022	10 ³ -10 ⁴	Good	Colorless	–
<i>Escherichia coli</i>	8739	<100	Growth (18-24 hours at 30-35°C)	Red to rose-red	+
<i>Escherichia coli</i>	8739	<100	Growth (18-72 hours at 35-37°C)	Red to rose-red	+

BBL™ MacConkey I Agar (prepared)

Inoculate and incubate at 35 ± 2°C for 18-24 hours. Incubate *E. coli* ATCC 8739 at 30-35°C for 18-72 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	29212	10 ⁴ -10 ⁵	Partial to complete inhibition	–	–
<i>Escherichia coli</i>	25922	10 ³ -10 ⁴	Good	Red to rose-red	+
<i>Pseudomonas aeruginosa</i>	10145	10 ³ -10 ⁴	Good	Greenish yellow	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	10 ³ -10 ⁴	Good	No reaction	–
<i>Shigella dysenteriae</i>	9361	10 ³ -10 ⁴	Good	No reaction	–
<i>Escherichia coli</i>	8739	10-100	Growth	Red to rose-red	+

BBL™ MacConkey Agar without Crystal Violet


Prepare the medium per label directions. Inoculate and incubate at 35 ± 2°C for 18-24 hours and up to 48 hours if necessary (up to 11 days for *M. fortuitum*).

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	29212	10 ³ -10 ⁴	Good	Rose red	–
<i>Escherichia coli</i>	25922	10 ³ -10 ⁴	Good	Pink to rose red	–
<i>Mycobacterium fortuitum</i>	6841	10 ³ -10 ⁴	Good	Rose red	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	10 ³ -10 ⁴	Good	Colorless	–
<i>Staphylococcus aureus</i>	25923	10 ³ -10 ⁴	Good	Pink to rose red	–

BBL™ MacConkey Agar without Crystal Violet or Salt

Prepare the medium per label directions. Inoculate and incubate at 35 ± 2°C for 18-24 hours and up to 48 hours if necessary (up to 11 days for *M. fortuitum*).

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY	COLONY COLOR	BILE PPT.
<i>Enterococcus faecalis</i>	29212	10 ³ -10 ⁴	Good	Rose red	–
<i>Escherichia coli</i>	25922	10 ³ -10 ⁴	Good	Pink to rose red	–
<i>Proteus mirabilis</i>	12453	10 ³ -10 ⁴	Good	Colorless, no swarming	–
<i>Salmonella enterica</i> subsp. <i>enterica</i> serotype Typhimurium	14028	10 ³ -10 ⁴	Good	Colorless	–



Summary and Explanation

MacConkey Agar is based on the bile salt-neutral red-lactose agar of MacConkey.¹²

The original MacConkey medium was used to differentiate strains of *Salmonella typhosa* from members of the coliform group. Formula modifications improved the growth of *Shigella* and *Salmonella* strains. These modifications included the addition of 0.5% sodium chloride, decreased agar content, and altered bile salts and neutral red concentrations. The formula improvements gave improved differential reactions between these enteric pathogens and the coliform group.

MacConkey Agar contains crystal violet and bile salts that inhibit gram-positive organisms and allow gram-negative organisms to grow. Isolated colonies of coliform bacteria are brick red in color and may be surrounded by a zone of precipitated bile. This bile precipitate is due to a local pH drop around the colony due to lactose fermentation. Colonies that do not ferment lactose (such as typhoid, paratyphoid and dysentery bacilli) remain colorless. When lactose nonfermenters grow in proximity to coliform colonies, the surrounding medium appears as cleared areas. MacConkey Agar is listed as one of the recommended media for the isolation of *E. coli* from nonsterile pharmaceutical products.⁹

MacConkey Agar Base is prepared without added carbohydrates, which permits their addition either individually or in combination. It is recommended that carbohydrates such as sucrose or lactose be added in a concentration of 1% to the basal medium.

MacConkey Agar without Crystal Violet is a differential medium that is less selective than MacConkey Agar. The lack of crystal violet permits the growth of *Staphylococcus* and *Enterococcus*.

Staphylococci produce pale pink to red colonies and enterococci produce compact tiny red colonies either on or beneath the surface of the medium. The medium is used also to separate *Mycobacterium fortuitum* and *M. chelonae* from other rapidly growing mycobacteria.^{13,14}

MacConkey Agar without Crystal Violet or Salt and MacConkey Agar without Salt (which also lacks crystal violet) are differential media used for isolating and cultivating gram-negative enteric organisms and gram-positive cocci from waters, feces and other sources suspected of containing these organisms, as well as limiting the swarming of *Proteus* species.

Principles of the Procedure

Peptones are sources of nitrogen and other nutrients. Yeast extract is a source of trace elements, vitamins, amino acids and carbon. Lactose is a fermentable carbohydrate. When lactose is fermented, a local pH drop around the colony causes a color change in the pH indicator (neutral red) and bile precipitation. Bile salts, bile salts no. 3, oxgall and crystal violet are selective agents that inhibit growth of gram-positive organisms. Sodium chloride maintains osmotic balance in the medium. Magnesium sulfate is a source of divalent cations. Agar is the solidifying agent.

Formulae

Difco™ MacConkey Agar

Approximate Formula* Per Liter	
Pancreatic Digest of Gelatin	17.0 g
Peptones (meat and casein).....	3.0 g
Lactose	10.0 g
Bile Salts No. 3	1.5 g
Sodium Chloride	5.0 g
Agar	13.5 g
Neutral Red.....	0.03 g
Crystal Violet.....	1.0 mg

Difco™ MacConkey Agar Base

Consists of the same ingredients without the lactose.

BBL™ MacConkey Agar

Approximate Formula* Per Liter	
Pancreatic Digest of Gelatin	17.0 g
Peptones (meat and casein).....	3.0 g
Lactose	10.0 g
Bile Salts	1.5 g
Sodium Chloride	5.0 g
Agar	13.5 g
Neutral Red.....	0.03 g
Crystal Violet.....	1.0 mg

Difco™ MacConkey Agar without Crystal Violet

Approximate Formula* Per Liter	
Peptone	20.0 g
Lactose	10.0 g
Bile Salts	5.0 g
Sodium Chloride	5.0 g
Agar	12.0 g
Neutral Red.....	0.05 g

BBL™ MacConkey Agar without Crystal Violet

Approximate Formula* Per Liter	
Pancreatic Digest of Casein	10.0 g
Peptic Digest of Animal Tissue.....	10.0 g
Lactose	10.0 g
Bile Salts	5.0 g
Sodium Chloride	5.0 g
Agar	12.0 g
Neutral Red.....	0.05 g

Difco™ MacConkey Agar without Salt

Approximate Formula* Per Liter	
Peptone	20.0 g
Lactose	10.0 g
Bile Salts	5.0 g
Agar	12.0 g
Neutral Red.....	75.0 mg

BBL™ MacConkey Agar without Crystal Violet or Salt

Approximate Formula* Per Liter	
Pancreatic Digest of Gelatin	10.0 g
Yeast Extract	10.0 g
Lactose	10.0 g
Oxgall	5.0 g
Magnesium Sulfate	0.2 g
Agar	12.0 g
Neutral Red.....	75.0 mg

*Adjusted and/or supplemented as required to meet performance criteria.

Directions for Preparation from Dehydrated Product

- Suspend the powder in 1 L of purified water:
 - Difco™ MacConkey Agar – 50 g;
 - BBL™ MacConkey Agar – 50 g;
 - Difco™ MacConkey Agar Base – 40 g;
 - Difco™ MacConkey Agar without Crystal Violet – 52 g;
 - BBL™ MacConkey Agar without Crystal Violet – 52 g;
 - BBL™ MacConkey Agar without Crystal Violet or Salt – 47.3 g;
 - Difco™ MacConkey Agar without Salt – 47 g.
 Mix thoroughly.
 - Heat with frequent agitation and boil for 1 minute to completely dissolve the powder.
 - Autoclave at 121°C for 15 minutes.
- NOTE: If MacConkey Agar Base is to be used within 12 hours, omit autoclaving and gently boil medium for 5 minutes. Add 1% carbohydrate before or after autoclaving, depending upon heat lability. The surface of MacConkey agars without salt should be thoroughly air-dried prior to inoculation.
- Test samples of the finished product for performance using stable, typical control cultures.

Sample Collection and Handling

For clinical specimens, refer to laboratory procedures for details on specimen collection and handling.¹⁻³

For food or dairy samples, follow appropriate standard methods for details on sample collection and preparation according to sample type and geographic location.⁴⁻⁷

For cosmetics, water, or other industrial samples, follow appropriate standard methods for details on sample collection and preparation according to sample type and geographic location.⁶⁻¹¹

For pharmaceutical samples, refer to *USP* General Chapter <62> for details on the examination of nonsterile products and tests for isolating *E. coli* using MacConkey Agar.⁹

Procedure

Refer to appropriate standard references for details on test methods to obtain isolated colonies from specimens or samples using MacConkey Agar.¹⁻¹¹ Incubate plates for 18-72 hours at 35 ± 2°C under appropriate atmospheric conditions, or as instructed in the standard reference.¹⁻¹¹

Expected Results

Lactose-fermenting organisms grow as pink to brick-red colonies with or without a zone of precipitated bile. Lactose-nonfermenting organisms grow as colorless or clear colonies.

Swarming by *Proteus* spp. is reduced on MacConkey agars without salt.

On MacConkey Agar without Crystal Violet and MacConkey agars without salt, staphylococci produce pale pink to red colonies and enterococci produce tiny red colonies; these organisms are inhibited on MacConkey Agar.

On MacConkey Agar without Crystal Violet, potentially pathogenic rapid growers of the *M. fortuitum* complex usually grow in 5-11 days, while the commonly saprophytic species are inhibited.^{3,13}

On MacConkey agars without salt, the swarming of *Proteus* is reduced.

Limitations of the Procedure

- Although MacConkey media are selective primarily for gram-negative enteric bacilli, biochemical and, if indicated, serological testing using pure cultures are recommended for complete identification. Consult appropriate references for further information.^{1,3}
- Incubation of MacConkey Agar plates under increased CO₂ has been reported to reduce the growth and recovery of a number of strains of gram-negative bacilli.¹⁴
- Some strains of *M. smegmatis* from humans may grow on MacConkey Agar without Crystal Violet, but these strains can be differentiated from *M. fortuitum* complex by the 3-day arylsulfatase test.⁹

References

- Murray, Baron, Jorgensen, Landry and Pfäller (eds.). 2007. Manual of clinical microbiology, 9th ed. American Society for Microbiology, Washington, D.C.
- Forbes, Sahm and Weissfeld. 2007. Bailey & Scott's diagnostic microbiology, 12th ed. Mosby Elsevier, St. Louis, Mo.
- Isenberg and Garcia (eds.). 2004 (update, 2007). Clinical microbiology procedures handbook, 2nd ed. American Society for Microbiology, Washington, D.C.
- Wehr and Frank (eds.). 2004. Standard methods for the examination of dairy products, 17th ed. American Public Health Association, Washington, D.C.
- Downes and Ito (ed.). 2001. Compendium of methods for the microbiological examination of foods, 4th ed. American Public Health Association, Washington, D.C.
- U.S. Food and Drug Administration. 2001. Bacteriological analytical manual, online (25 Sept 2008). AOAC International, Gaithersburg, Md.
- Horwitz (ed.). 2007. Official methods of analysis of AOAC International, 18th ed., online. AOAC International, Gaithersburg, Md.
- Eaton, Rice and Baird (eds.). 2005. Standard methods for the examination of water and wastewater, 21st ed., online. American Public Health Association, Washington, D.C.
- United States Pharmacopeial Convention, Inc. 2008. The United States pharmacopeia 31/The national formulary 26, Supp. 1, 8-1-08, online. United States Pharmacopeial Convention, Inc., Rockville, Md.

10. European Directorate for the Quality of Medicines and Healthcare. 2008. The European pharmacopoeia, 6th ed., Supp. 1, 4-1-2008, online. European Directorate for the Quality of Medicines and Healthcare, Council of Europe, 226 Avenue de Colmar BP907, F-67029 Strasbourg Cedex 1, France.
11. Japanese Ministry of Health, Labour and Welfare. 2006. The Japanese pharmacopoeia, 15th ed., online. Japanese Ministry of Health, Labour and Welfare.
12. MacConkey. 1905. J. Hyg. 5:333.
13. Kent and Kubica. 1985. Public health mycobacteriology: a guide for the level III laboratory. USDHHS, Centers for Disease Control, Atlanta, Ga.
14. Mazura-Reetz, Neblett and Galperin. 1979. Abstr. C179, p. 339. Abstr. Annu. Meet. American Society for Microbiology, 1979.

Availability

Difco™ MacConkey Agar

AOAC BAM BS12 CCAM CMPH2 COMPF EP JP MCM9
SMD SMWW USP

Cat. No. 212123 Dehydrated – 500 g[†]
212122 Dehydrated – 2 kg[†]
275300 Dehydrated – 10 kg[†]

BBL™ MacConkey Agar

AOAC BAM BS12 CCAM CMPH2 COMPF EP JP MCM9
SMD SMWW USP

Cat. No. 211387 Dehydrated – 500 g[†]
211390 Dehydrated – 5 lb (2.3 kg)[†]
211391 Dehydrated – 25 lb (11.3 kg)[†]

BBL™ MacConkey I Agar

AOAC BAM BS12 CCAM CMPH2 COMPF EP JP MCM9
SMD SMWW USP

United States and Canada

Cat. No. 215197 Prepared Plates – Pkg. of 20*[†]
297064 Prepared Plates – Ctn. of 100*[†]

Difco™ MacConkey Agar Base

Cat. No. 281810 Dehydrated – 500 g

Difco™ MacConkey Agar without Crystal Violet

Cat. No. 247010 Dehydrated – 500 g

BBL™ MacConkey Agar without Crystal Violet

Cat. No. 211393 Dehydrated – 500 g

Europe

Cat. No. 256008 Prepared Plates – Pkg. of 20*

BBL™ MacConkey Agar without Crystal Violet or Salt

Cat. No. 294584 Dehydrated – 500 g

Difco™ MacConkey Agar without Salt

Cat. No. 233120 Dehydrated – 500 g
233110 Dehydrated – 10 kg

Europe

Cat. No. 256009 Prepared Plates – Pkg. of 20*
257286 Prepared Plates – Ctn. of 120*

* Store at 2-8°C.

[†] QC testing performed according to USPIEPIIP performance specifications.

MacConkey II Agar • MacConkey II Agar with MUG

Intended Use

MacConkey II Agar is a slightly selective and differential medium for the detection of coliform organisms and enteric pathogens.

MacConkey II Agar with MUG is used for the presumptive identification of *Escherichia coli*.

MacConkey II Agar meets *United States Pharmacopeia (USP)* performance specifications.

Summary and Explanation

The BBL™ MacConkey II Agar formulation was made available in 1983. It was specially designed to improve the inhibition of swarming *Proteus* species, to achieve more definitive differentiation of lactose fermenters and nonfermenters, and for the promotion of superior growth of enteric pathogens.

Trepeta and Edberg¹ modified MacConkey Agar by the incorporation of MUG (4-methylumbelliferyl-β-D-glucuronide). The resulting medium allowed the authors to presumptively identify *E. coli* from the primary plating medium within 5 minutes.

Principles of the Procedure

MacConkey II Agar is a selective and differential medium. It is only slightly selective since the concentration of bile salts, which inhibit gram-positive microorganisms, is low in comparison with other enteric plating media. Crystal violet also is included in the medium to inhibit the growth of gram-positive bacteria, especially enterococci and staphylococci.

Differentiation of enteric microorganisms is achieved by the combination of lactose and the neutral red indicator. Colorless or pink to red colonies are produced depending upon the ability of the isolate to ferment the carbohydrate.

Most strains (96-97%) of *E. coli* produce β-D-glucuronidase.² The enzyme hydrolyzes MUG to yield 4-methylumbelliferone, a compound that fluoresces under long-wave (366 nm) UV light. The addition of MUG to the formulation allows β-D-glucuronidase-positive strains of *E. coli* to fluoresce blue-green when examined under UV light.

BBL MacConkey II Agar with MUG contains 0.1 g of MUG per liter of MacConkey II Agar.

Formula

BBL™ MacConkey II Agar

Approximate Formula* Per Liter

Pancreatic Digest of Gelatin	17.0	g
Pancreatic Digest of Casein	1.5	g
Peptic Digest of Animal Tissue.....	1.5	g
Lactose	10.0	g
Bile Salts	1.5	g
Sodium Chloride	5.0	g
Agar	13.5	g
Neutral Red.....	0.03	g
Crystal Violet.....	1.0	mg

*Adjusted and/or supplemented as required to meet performance criteria.