

Test report No. shd1118

EVALUATION OF BACTERICIDAL ACTIVITIES OF A DISINFECTANTS AND ANTISEPTICS USED IN THE MEDICAL AREA (EN 13727)

Name of the product: Chemisept GEL  
Batch number: 198251017  
Order number: 17030  
Manufacturer: Chemi-Pharm Ltd  
Client, representative: Chemi-Pharm Ltd., Põllu 132, Tallinn, 10917, ESTONIA  
Maris Millner, +372-51-77-090  
Date of delivery: 15.01.2018  
Test material conditions: No specific features, sample in the manufacturers tare  
Storage conditions: In room temperature, dark  
Active substance – conc.: Ethyl alcohol – 72.5 % wt; isopropyl alcohol – 7.5 % wt  
Appearance of the product: Transparent liquid  
Test concentration: Ready to use  
Contact time: 15sec, 30 sec  
Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions; 1,5g/l bovine albumin = clean conditions  
Neutralizer: -  
Rinsing liquid: Tryptone 1 g/l + NaCl, 9 g/l  
Test organisms: *Pseudomonas aeruginosa* ATCC 15442  
*Staphylococcus aureus* ATCC 6538  
*Enterococcus hirae* ATCC 10541  
*Escherichia coli* K12, NTCT 10538  
*Staphylococcus aureus* MRSA ATCC 33592  
*Enterococcus faecium* VRE ATCC 700221  
Testing method: EVS-EN 13727:2012+A2:2015  
Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity in the medical area - Test method and requirements (phase 2, step 1)  
Testing date: 17.02.2018 – 19.02.2018  
Results: look appendix 1-7



Diana Kaare, MSc  
Head of laboratory, microbiologist  
Date of test report: 21.02.2018

Appendix 1

TEST RESULTS (bactericidal suspension test)

EVS-EN 13727:2012+A2:2015; Phase 2, step 1;  
 Membrane filtration method;  
 Rinsing liquid: tryptone, 1 g/l+ NaCl, 9 g/l;  
 Test organism: *Staphylococcus aureus* ATCC 6538;  
 Test temperature: +20° C; Incubation temperature: +37 °C  
 Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions;  
 1,5g/l bovine albumin = clean conditions  
 Nordic Tersus Laboratory LLC.;  
 Date of test: 17.02.2018 – 19.02.2018  
 Responsible person: Diana Kaare

Validation and controls

Dirty and clean conditions

Validation suspension $N_{vo}$			Experimental conditions control (A)			Filtration control (B)			Method validation (C)		
$V_{C1}$	83	$\bar{x} = 86.5$	$V_{C1}$	72	$\bar{x} = 68.5$	$V_{C1}$	68	$\bar{x} = 71.5$	$V_{C1}$	81	$\bar{x} = 77$
$V_{C2}$	90		$V_{C2}$	65		$V_{C2}$	75		$V_{C2}$	73	
$30 \leq \bar{x} N_{vo} \leq 160$ ? yes X; no <input type="checkbox"/>			$\bar{x} A$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} B$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} C$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>		

Test suspension and test

Testsuspension:	$N$	$V_{C1}$	$V_{C2}$	$\bar{x}_{wm} = 2.45 \times 10^9$ ; $\log N = 9.39$
$N$ and $N_o$	$10^{-7}$	257	224	$N_o = N/100$ ; $\log N_o = 7,39$
	$10^{-8}$	28	30	$7,17 \leq \log N_o \leq 7,70$ ; yes X; no <input type="checkbox"/>

Experimental results

Concentration of the product %	Dilution step	$V_{C1}$	$V_{C2}$	Na (= $\bar{x}$ *10)	log Na	logR	Contact time	Conditions
Ready to use	-	<14	<14	<140	<2.15	>5.24	15 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.24	15 sec	clean
Ready to use	-	<14	<14	<140	<2.15	>5.24	30 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.24	30 sec	clean

Explanations:

$V_C$  = count per ml (one plate or more)

$\bar{x}$  = average of  $V_{C1}$  and  $V_{C2}$  (1. + 2. duplicate)

$N$  = cfu/ml microbes in testsuspension

$N_0$  = cfu/ml at the start of the contact time (t=0)

$N_{vo}$  = cfu/ml in the validation suspension (t=0)

$N_a$  = surviving microbes after the test

R = reduction factor ( $R = N_0 / N_a$ ;  $\text{LogR} = \text{Log}N_0 - \text{Log}N_a$ )

Appendix 2

TEST RESULTS (bactericidal suspension test)

EVS-EN 13727:2012+A2:2015; Phase 2, step 1;  
 Membrane filtration method;  
 Rinsing liquid: tryptone, 1 g/l+ NaCl, 9 g/l;  
 Test organism: *Enterococcus hirae* ATCC 10541;  
 Test temperature: +20° C; Incubation temperature: +37 °C  
 Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions;  
 1,5g/l bovine albumin = clean conditions  
 Nordic Tersus Laboratory LLC.;  
 Date of test: 17.02.2018 – 19.02.2018  
 Responsible person: Diana Kaare

Validation and controls

Dirty and clean conditions

Validation suspension $N_{vo}$			Experimental conditions control (A)			Filtration control (B)			Method validation (C)		
$V_{C1}$	62	$\bar{x} = 66$	$V_{C1}$	47	$\bar{x} = 46$	$V_{C1}$	52	$\bar{x} = 53,5$	$V_{C1}$	59	$\bar{x} = 61$
$V_{C2}$	70		$V_{C2}$	45		$V_{C2}$	55		$V_{C2}$	53	
$30 \leq \bar{x} N_{vo} \leq 160$ ? yes X; no <input type="checkbox"/>			$\bar{x} A$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} B$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} C$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>		

Test suspension and test

Testsuspension:	$N$	$V_{C1}$	$V_{C2}$	$\bar{x}_{wm} = 2.15 \times 10^9$ ; $\log N = 9.33$
$N$ and $N_0$	$10^{-7}$	203	224	$N_0 = N/100$ ; $\log N_0 = 7,33$
	$10^{-8}$	22	25	$7,17 \leq \log N_0 \leq 7,70$ ; yes X; no <input type="checkbox"/>

Experimental results

Concentration of the product %	Dilution step	$V_{C1}$	$V_{C2}$	Na (= $\bar{x}$ *10)	log Na	logR	Contact time	Conditions
Ready to use	-	<14	<14	<140	<2.15	>5.18	15 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.18	15 sec	clean
Ready to use	-	<14	<14	<140	<2.15	>5.18	30 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.18	30 sec	clean

Explanations:

$V_C$  = count per ml (one plate or more)

$\bar{x}$  = average of  $V_{C1}$  and  $V_{C2}$  (1. + 2. duplicate)

$N$  = cfu/ml microbes in testsuspension

$N_0$  = cfu/ml at the start of the contact time (t=0)

$N_{vo}$  = cfu/ml in the validation suspension (t=0)

$N_a$  = surviving microbes after the test

R = reduction factor (R=  $N_0/ N_a$ ; LogR=Log $N_0$  - Log  $N_a$ )

Appendix 3

**TEST RESULTS (bactericidal suspension test)**

EVS-EN 13727:2012+A2:2015; Phase 2, step 1;

Membrane filtration method;

Rinsing liquid: tryptone, 1 g/l+ NaCl, 9 g/l;

Test organism: *Pseudomonas aeruginosa* ATCC 15442

Test temperature: +20° C; Incubation temperature: +37 °C

Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions;

1,5g/l bovine albumin = clean conditions

Nordic Tersus Laboratory LLC.;

Date of test: 17.02.2018 – 19.02.2018

Responsible person: Diana Kaare

Validation and controls

Dirty and clean conditions

Validation suspension $N_{vo}$			Experimental conditions control (A)			Filtration control (B)			Method validation (C)		
$V_{C1}$	94	$\bar{x} = 87$	$V_{C1}$	71	$\bar{x} = 72.5$	$V_{C1}$	66	$\bar{x} = 68$	$V_{C1}$	75	$\bar{x} = 78.5$
$V_{C2}$	80		$V_{C2}$	74		$V_{C2}$	70		$V_{C2}$	82	
$30 \leq \bar{x} N_{vo} \leq 160$ ? yes X; no <input type="checkbox"/>			$\bar{x} A$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} B$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} C$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>		

Test suspension and test

Testsuspension:	$N$	$V_{C1}$	$V_{C2}$	$\bar{x}_{wm} = 2.74 \times 10^9$ ; $\log N = 9.44$
$N$ and $N_0$	$10^{-7}$	284	266	$N_0 = N/100$ ; $\log N_0 = 7,44$
	$10^{-8}$	24	29	$7,17 \leq \log N_0 \leq 7,70$ ; yes X; no <input type="checkbox"/>

Experimental results

Concentration of the product %	Dilution step	$V_{C1}$	$V_{C2}$	Na (= $\bar{x}$ *10)	log Na	logR	Contact time	Conditions
Ready to use	-	<14	<14	<140	<2.15	>5.29	15 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.29	15 sec	clean
Ready to use	-	<14	<14	<140	<2.15	>5.29	30 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.29	30 sec	clean

Explanations:

$V_C$  = count per ml (one plate or more)

$\bar{x}$  = average of  $V_{C1}$  and  $V_{C2}$  (1. + 2. duplicate)

$N$  = cfu/ml microbes in testsuspension

$N_0$  = cfu/ml at the start of the contact time (t=0)

$N_{v0}$  = cfu/ml in the validation suspension (t=0)

$N_a$  = surviving microbes after the test

R = reduction factor ( $R = N_0 / N_a$ ;  $\text{LogR} = \text{Log}N_0 - \text{Log}N_a$ )

**TEST RESULTS (bactericidal suspension test)**

EVS-EN 13727:2012+A2:2015; Phase 2, step 1;

Membrane filtration method;

Rinsing liquid: tryptone, 1 g/l+ NaCl, 9 g/l;

Test organism: *Escherichia coli* K12, NTCT 10538

Test temperature: +20° C; Incubation temperature: +37 °C

Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions;

1,5g/l bovine albumin = clean conditions

Nordic Tersus Laboratory LLC.;

Date of test: 17.02.2018 – 19.02.2018

Responsible person: Diana Kaare

**Validation and controls**

**Dirty and clean conditions**

Validation suspension $N_{vo}$			Experimental conditions control (A)			Filtration control (B)			Method validation (C)		
$V_{C1}$	82	$\bar{x} = 78.5$	$V_{C1}$	68	$\bar{x} = 60$	$V_{C1}$	69	$\bar{x} = 69.5$	$V_{C1}$	70	$\bar{x} = 68.5$
$V_{C2}$	75		$V_{C2}$	52		$V_{C2}$	73		$V_{C2}$	67	
$30 \leq \bar{x} N_{vo} \leq 160$ ? yes X; no <input type="checkbox"/>			$\bar{x} A$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} B$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} C$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>		

**Test suspension and test**

Testsuspension:	$N$	$V_{C1}$	$V_{C2}$	$\bar{x}_{wm} = 2.48 \times 10^9$ ; $\log N = 9.39$ $N_0 = N/100$ ; $\log N_0 = 7.39$ $7,17 \leq \log N_0 \leq 7,70$ ; yes X; no <input type="checkbox"/>
$N$ and $N_0$	$10^{-7}$	241	257	
	$10^{-8}$	22	26	



Experimental results

Concentration of the product %	Dilution step	$V_{C1}$	$V_{C2}$	Na (= $\bar{x}$ *10)	log Na	logR	Contact time	Conditions
Ready to use	-	<14	<14	<140	<2.15	>5.24	15 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.24	15 sec	clean
Ready to use	-	<14	<14	<140	<2.15	>5.24	30 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.24	30 sec	clean

Explanations:

$V_C$  = count per ml (one plate or more)

$\bar{x}$  = average of  $V_{C1}$  and  $V_{C2}$  (1. + 2. duplicate)

$N$  = cfu/ml microbes in testsuspension

$N_0$  = cfu/ml at the start of the contact time (t=0)

$N_{vo}$  = cfu/ml in the validation suspension (t=0)

$N_a$  = surviving microbes after the test

$R$  = reduction factor ( $R = N_0 / N_a$ ;  $\text{Log}R = \text{Log}N_0 - \text{Log}N_a$ )

**TEST RESULTS (bactericidal suspension test)**

EVS-EN 13727:2012+A2:2015; Phase 2, step 1;

Membrane filtration method;

Rinsing liquid: tryptone, 1 g/l+ NaCl, 9 g/l;

Test organism: *Staphylococcus aureus* MRSA ATCC 33592

Test temperature: +20° C; Incubation temperature: +37 °C

Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions;

1,5g/l bovine albumin = clean conditions

Nordic Tersus Laboratory LLC.;

Date of test: 17.02.2018 – 19.02.2018

Responsible person: Diana Kaare

**Validation and controls**

**Dirty and clean conditions**

Validation suspension $N_{vo}$			Experimental conditions control (A)			Filtration control (B)			Method validation (C)		
$V_{C1}$	66	$\bar{x} = 61.5$	$V_{C1}$	42	$\bar{x} = 45.5$	$V_{C1}$	49	$\bar{x} = 47.5$	$V_{C1}$	52	$\bar{x} = 53$
$V_{C2}$	57		$V_{C2}$	49		$V_{C2}$	46		$V_{C2}$	54	
$30 \leq \bar{x} N_{vo} \leq 160$ ? yes X; no <input type="checkbox"/>			$\bar{x} A$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} B$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} C$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>		

**Test suspension and test**

Testsuspension:	$N$	$V_{C1}$	$V_{C2}$	$\bar{x}_{wm} = 1.99 \times 10^9$ ; $\log N = 9.30$ $N_0 = N/100$ ; $\log N_0 = 7.30$ $7,17 \leq \log N_0 \leq 7,70$ ; yes X; no <input type="checkbox"/>
$N$ and $N_0$	$10^{-7}$	204	191	
	$10^{-8}$	18	24	

Experimental results

Concentration of the product %	Dilution step	$V_{C1}$	$V_{C2}$	Na (= $\bar{x}$ *10)	log Na	logR	Contact time	Conditions
Ready to use	-	<14	<14	<140	<2.15	>5.15	15 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.15	15 sec	clean
Ready to use	-	<14	<14	<140	<2.15	>5.15	30 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.15	30 sec	clean

Explanations:

$V_C$  = count per ml (one plate or more)

$\bar{x}$  = average of  $V_{C1}$  and  $V_{C2}$  (1. + 2. duplicate)

$N$  = cfu/ml microbes in testsuspension

$N_0$  = cfu/ml at the start of the contact time (t=0)

$N_{vo}$  = cfu/ml in the validation suspension (t=0)

$Na$  = surviving microbes after the test

R = reduction factor (R=  $N_0/ Na$ ; LogR=Log $N_0$  - Log  $Na$ )

**TEST RESULTS (bactericidal suspension test)**

EVS-EN 13727:2012+A2:2015; Phase 2, step 1;

Membrane filtration method;

Rinsing liquid: tryptone, 1 g/l+ NaCl, 9 g/l;

Test organism: *Enterococcus faecium* VRE ATCC 700221

Test temperature: +20° C; Incubation temperature: +37 °C

Interfering substance: 15 g/l bovine albumin + 15 ml/l sheep blood erythrocytes = Dirty conditions;

1,5g/l bovine albumin = clean conditions

Nordic Tersus Laboratory LLC.;

Date of test: 17.02.2018 – 19.02.2018

Responsible person: Diana Kaare

**Validation and controls**

**Dirty and clean conditions**

Validation suspension $N_{vo}$			Experimental conditions control (A)			Filtration control (B)			Method validation (C)		
$V_{C1}$	62	$\bar{x} = 64$	$V_{C1}$	52	$\bar{x} = 54.5$	$V_{C1}$	43	$\bar{x} = 47$	$V_{C1}$	58	$\bar{x} = 59$
$V_{C2}$	66		$V_{C2}$	57		$V_{C2}$	51		$V_{C2}$	60	
$30 \leq \bar{x} N_{vo} \leq 160$ ? yes X; no <input type="checkbox"/>			$\bar{x} A$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} B$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>			$\bar{x} C$ is $\geq 0,5 \bar{x} N_{vo}$ ? yes X; no <input type="checkbox"/>		

**Test suspension and test**

Testsuspension:	$N$	$V_{C1}$	$V_{C2}$	$\bar{x}_{wm} = 1.88 \times 10^9$ ; $\log N = 9.27$
$N$ and $N_0$	$10^{-7}$	181	195	$N_0 = N/100$ ; $\log N_0 = 7.27$
	$10^{-8}$	22	18	$7,17 \leq \log N_0 \leq 7,70$ ; yes X; no <input type="checkbox"/>

Experimental results

Concentration of the product %	Dilution step	$V_{C1}$	$V_{C2}$	Na (= $\bar{x}$ *10)	log Na	logR	Contact time	Conditions
Ready to use	-	<14	<14	<140	<2.15	>5.12	15 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.12	15 sec	clean
Ready to use	-	<14	<14	<140	<2.15	>5.12	30 sec	dirty
Ready to use	-	<14	<14	<140	<2.15	>5.12	30 sec	clean

Explanations:

$V_C$  = count per ml (one plate or more)

$\bar{x}$  = average of  $V_{C1}$  and  $V_{C2}$  (1. + 2. duplicate)

$N$  = cfu/ml microbes in testsuspension

$N_0$  = cfu/ml at the start of the contact time (t=0)

$N_{vo}$  = cfu/ml in the validation suspension (t=0)

$Na$  = surviving microbes after the test

R = reduction factor ( $R = N_0 / Na$ ;  $\text{LogR} = \text{Log}N_0 - \text{Log}Na$ )

Appendix 7

Interpretation:

The product for surgical handrub Chemisept GEL (batch no. 198251017) was tested according to the test method EVS-EN 13727:2012+A2:2015. The test was performed at  $20\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$ , under clean and dirty conditions with the contact times of 15 sec and 30 sec. The membrane filtration method was used for testing the products' effectiveness against the reference strains: *Pseudomonas aeruginosa* ATCC 15442, *Enterococcus hirae* ATCC 10541, *Staphylococcus aureus* ATCC 6538, Escherichia coli K12 NTCT 10538, Staphylococcus aureus MRSA ATCC 33592 and *Enterococcus faecium* VRE ATCC 700221. Under clean and dirty conditions the tested product was effective against all the reference strains within 15 sec and 30 sec of contact times.

Conclusion:

The surviving count of bacterial reference strains showed at least 5lg reduction meaning that under clean and dirty conditions the ready to use product Chemisept GEL has a bactericidal effect in case of hand disinfection within 15 sec.



Diana Kaare, MSc

Head of laboratory, microbiologist