



## V Factor Discs

DD021

Used for the presumptive identification of *Haemophilus* species on the basis of their requirements for X or V factors or both.

### Directions

Inoculate the surface of a Blood Agar (M073) plate or Brain Heart Infusion Agar (M211) plate with the test organisms by either streaking or surface spreading. Aseptically place the X (DD020), V (DD021) and X+V (DD022) factor discs on the plate, in the following positions:

Disc Position on the Agar plate

X factor disc 12 O' clock

V factor disc 4 O' clock

X+V factor disc 8 O' clock

Incubate the plates at 35 - 37°C for 24 - 48 hours. Observe for the growth in the neighbourhood of the discs.

### Principle And Interpretation

Both X and V factors are growth factors that are essential for certain organisms like *Haemophilus* species and also enhance growth of organisms like *Neisseria* species.

V-factor discs are the sterile filter paper discs impregnated with growth factor V which are used for differentiating *Haemophilus* species in conjunction of X factor & X+V factor discs. *Bordetella* and *Haemophilus* species can also be identified on the basis of the requirement of X and V growth factors in the basal medium.

The X factor (hemin) and V factor (Coenzyme- Nicotinamide adenine dinucleotide NAD+) are impregnated on the sterile filter paper discs of diameter 6 mm.

The test organism requiring X factor alone, grows only in the vicinities of X and X+V factor discs. Those which require V factor alone grow in the vicinities of V and X+V factor discs. If both X and V factors are required, then the organism will grow only in the vicinity of the X+V factor discs. This satellite growth is seen around the disc promoting growth (1).

### Quality Control

#### Appearance

Filter paper discs of 6 mm diameter bearing letters "V" in continuous printing style.

#### Cultural response

Cultural characteristics observed on Brain Heart Infusion Agar (M211) or Blood Agar Base (M073) after an incubation of 24-48 hours at 35-37°C.

Organism	Growth with V factor	Growth without growth factor
<i>Bordetella pertussis</i> ATCC 8467	Positive(initial isolation on Bordet Gengou Agar (M175))	Positive(initial isolation on Bordet Gengou Agar (M175))
<i>Haemophilus influenzae</i> ATCC 35056	Negative	Negative
<i>Haemophilus parainfluenzae</i> ATCC 7901	Positive	Negative

---

<i>Haemophilus</i> <i>haemoglobinophilus</i> ATCC19416	Negative	Negative
<i>Haemophilus ducreyi</i>	Negative	Negative

### Storage and Shelf Life

Store below -10°C. Use before the expiry date on the label.

### Reference

1.Murray PR, Baron EJ, Jorgensen J.H., Pfaller M A, Tenover F.C., Tenover P.C.(Eds.),8th ed, 2003, Manual of Clinical Microbiology, ASM, Washington D.C.

#### Note:

Use known strains of *Haemophilus influenzae* to monitor the performance of the differentiation discs and the medium.

Do not use too heavy suspension of the test organisms as X or V factor carryover from the primary growth medium may take place

Revision : 1 / 2011



#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.