



# Technical Data

## Fraser Selective Supplement

FD125I

Recommended for selective isolation and enumeration of *Listeria monocytogenes* from food, animal feeds etc.

### Composition

Per vial sufficient for 500 ml / 1000 ml medium

#### \*Ingredients

Acriflavin hydrochloride

Nalidixic acid

#### Concentration

12.500mg

10mg

### Directions:

Rehydrate the contents of 1 vial aseptically with 10 ml of sterile distilled water. Mix well. Aseptically add 1 vial to 1000 ml sterile, cooled (45-50°C) Fraser Broth Base [M1327](#) / Fraser HiVeg™ Broth Base [MV1327](#) / Fraser Enrichment Broth Base [M1083R](#) for primary enrichment or to 500 ml sterile, Fraser Broth Base [M1327](#) / Fraser HiVeg™ Broth Base [MV1327](#) / Fraser Enrichment Broth Base [M1083R](#) for secondary enrichment. Mix well and dispense as desired.

### Storage and Shelf Life

Store at 2-8°C. Use before the expiry date on the label.

Revision : 1 / 2012

\* Not For Medicinal Use

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.