

SABOURAUD DEXTROSE (2%) AGAR

A non-selective medium for the cultivation and isolation of pathogenic and non-pathogenic fungi, particularly dermatophytes.

Dehydrated media	
Code number:	500 g: SDD20500, 5 kg: SDD25000
Colour:	Yellowish
Appearance:	Homogeneous hygroscopic powder
pH before autoclaving (25 °C):	6,2 – 6,6

Direction: Suspend **45 g** in one litre of distilled water and heat with frequent agitation until the medium boils well. Sterilise by autoclaving at 121 °C for 15 minutes. Mix well and pour into Petri-dishes or tubes (cooling in slanted position).

Prepared media	
Bottled media:	100 ml: SDD30100, 500 ml: SDD30500
Plated media:	55 mm: SDD50055, 90 mm: SDD50090
Tubed media:	100 x 15 mm: SDD40005 (5 ml, slant)
Colour:	Yellowish
pH (25 °C):	6,3 – 6,5

Direction: Dispense the melted bottled media aseptically into sterile Petri-dishes or tubes (cooling in slanted position). Media in Petri-dishes and tubes are ready to use.

FORMULA in g/l

Peptones	10
Glucose	20
Agar	15

Note: The typical formula can be adjusted to obtain optimal performance.

Storage conditions: Store the dehydrated media tightly closed in a dry place at room temperature. Store the bottled media protected from light at room temperature. Store the plated and tubed media protected from light at 2-8 °C. Use before the expiry date on the label.

Quality control:

Test strains	Incubation temp: 25 °C	Growth	Incubation time: 48 h
<i>Candida albicans</i>	ATCC 10231	Good	
<i>Aspergillus brasiliensis</i>	ATCC 16404	Good	

References: Emmons et al. (1977) Medical Mycology

In vitro diagnostic – for professional use only!