

Dehydrating and rehydrating agents

Alcoholic solutions and absolute alcohol are ideal for histocytology and cytology processing, storing and transporting samples. They are also used for dehydration and rehydration of tested samples.

Histanol 50

Denaturated 50% ethyl alcohol.

1000 mL	H50-1L
5000 mL	H50-5L
10000 mL	H50-10L



Histanol 70

Denaturated 70% ethyl alcohol.

1000 mL	H70-1L
5000 mL	H70-5L
10000 mL	H70-10L



Histanol 80

Denaturated 80% ethyl alcohol.

1000 mL	H80-1L
5000 mL	H80-5L
10000 mL	H80-10L



Histanol 95

Denaturated 95% ethyl alcohol.

1000 mL	H95-1L
5000 mL	H95-5L
10000 mL	H95-10L



Histanol 96

Denaturated 96% ethyl alcohol.

1000 mL	H96-1L
5000 mL	H96-5L
10000 mL	H96-10L



Histanol 100

Denaturated 100% ethyl alcohol.

1000 mL	H100-1L
5000 mL	H100-5L
10000 mL	H100-10L



Histanol EP 95

95% mixture of denaturated ethyl alcohol and isopropyl alcohol.

1000 mL	HEP95-1L
5000 mL	HEP95-5L
10000 mL	HEP95-10L



Histanol EP 100

100% mixture of denaturated ethyl alcohol and isopropyl alcohol.

1000 mL	HEP100-1L
5000 mL	HEP100-5L
10000 mL	HEP100-10L



Histanol EP 9010

Alcoholic solvent substituting standard ethanol series in processing, dewaxing and dehydrating.

1000 mL	HEP91-1L
5000 mL	HEP91-5L
10000 mL	HEP91-10L



Histanol IP

Isopropyl alcohol for use in microscopy.

1000 mL	HIP-1L
5000 mL	HIP-5L
10000 mL	HIP-10L



Histanol M

Methyl alcohol for use in microscopy.

1000 mL	HM-1L
5000 mL	HM-5L
10000 mL	HM-10L



Histanol G

Glycerol for use in microscopy.

1000 mL	HG-1L
5000 mL	HG-5L
10000 mL	HG-10L

