

Pacric Set for cricothyrotomy

Pacric Set for cricothyrotomy



 \oplus



Packed in, rigid, single use blister pack to protect it from deformation and damage.

We offer following sets:

- sterile and ready to use, reference number 30-0001
- non-sterile, for training purposes only, reference number 30-0001N

in obsterics; 1:50-100 Emergency/ICU)[1]

[1] Cook T et al. Complications and failure of airw management. BJA 2012;109:i68-i89

Picture instruction of a cricothyrotomy procedure attached to each set





We also offer cricothyrotomy trainer for development of surgical cricothyrotomy skills. Each set contains:

- 1 neck base,
- 1 simulated skin,
- 1 simulated thyroid cartilage insert,
- 3 simulated thyroid membranes.

Individual components can be purchased separately. Complete set reference number: CR-TR-1.

Based on DAS 2015 guidelines

Guideliness of Difficult Airway Society (DAS 2015) Plan D: Emergency Front of Neck Access - surgical cricothyroidotomy.

> Ready for use when CICO declared. "Can't intubate, can't oxygenate"



SUMI spółka z ograniczoną odpowiedzialnością sp. k. Drobiarska 35, 05-070 Sulejówek, Poland e-mail: sumi@sumi.com.pl tel. +48 22 783 30 23 fax. +48 22 783 30 25



 \oplus

PaCric® **Set for cricothyrotomy**

According to DAS 2015 guildelines, in cases where it is not possible to:

- intubate patient,
- oxygenate patient,

- graduation marks,

- length adapted to cricothyrotomy procedure

- stop the procedure,
- it is necessary to move to Plan D, surgical cricothyrotomy procedure.

In anesthesiologist practice cicothyrotomy procedure happens extremely rarely. That is why it is important to be well acquainted with the procedure and necessary components.

PaCric set for cricothyrotomy is designed thinking of:

- -faster,
- safer,
- creating of emergency way for patient ventilation.

Tracheostomy tube with cuff, size 6,0 - reinforced - prevents tube kinking, - adjustable flange - provides tube length **Based on DAS** regulation, - Magil type - minimizes trauma during 2015 introduction. quideliness effective length - 15cm, flexible and arced - easier adaptation to patient **Cotton tape for** tracheostomy tube - pure, delicate cotton, easy to adjust to any neck size, minimizes skin irritations Syringe 10 ml Introducer - angled tip, bougie type, Scalpel, size 10 - atraumatic tip, - very flexible, shape retaining, - smooth surface minimizes resistance during

EMERGENCY LIFE SAVING PROCEDURE



Following cricothyrotomy procedure is recommended:



impalpable cricothyroid membrane



- Stand on the patient left-hand side.

incision, caudad to cephalad.

- Extend head pillow under shoulders.
 Tension skin using the left hand.
- Make an 8-10 cm midline vertical skin
- Using fingers of both hands separate tissues and identify cricothyroid membrane.
- Follow instructions from point 3 on the

PaCric cricothyrotomy set is used during 4-step surgical cricothyrotomy. Set utilizes medical devices which are well known to anaesthesiologists. It is adapted and assembled in 1 ready to use set, that provides intuitive usage during CICO cases (can't intubate, can't oxygenate)




palpable cricothyroid membrane






- Stand on the patient left-hand side.
- Extend head, pillow under shoulders. Using left hand identify cricothyroid



- Stabilize larynx using your left hand. Make a transverse stab incision through the skin and cricothyroid membrane with
- the blade cutting edge towards you.



Turn scalpel through 90° so that the sharp edge points caudally (towards the feet).



- Hold the scalpel with your left hand. Using right hand slide the tip of the
- introducer down the side of the scalpel blade, on 10-15cm caudally.
- Withdraw scalpel.



- Railroad tube over introducer. Withdraw introducer, inflate cuff and
- confirm correct tube placement.
- Secure the tube.



Instructions for use is provided with each set. It can also be download from our website www.sumi.com.pl, bookmark: Products, Set for cricothyrotomy