

Azure II Eosinate

Azure II eosinate. For preparation of Romanowsky stains, made by mixing equal ratios of Azure B, Methylene Blue and Eosin Y stains.

25 g	A2E-P-25
------	----------

Biebrich Scarlet, C.I. 26905

Croceine Scarlet, Ponceau B, Ponceau BS, BSC certified stain. For Masson trichrome staining and a component of Shorr's solution.

25 g	BS-P-25
------	---------

Bismarck Brown Y, C.I. 21000

Bismarck Brown Y, Vesuvine BA, BSC certified. For use in histopathology and cytology.

25 g	BB-P-25
------	---------

Brilliant Cresyl Blue, C.I. 51010

For use in hematology (for reticulocyte staining) and histology (for acid mucopolysaccharides).

10 g	BCB-P-10
25 g	BCB-P-25

Brilliant Green Certified, C.I. 42040

Malachite Green G, Basic Green 1, Emerald Green. For bacteriostatic use.

25 g	BG-P-25
------	---------

Carmine, C.I. 75470

Natural Red 4, Cochineal, Nacaral, carminic acid and aluminum compound. For staining mucins and glycogens.

5 g	CAR-P-5
10 g	CAR-P-10
25 g	CAR-P-25

Carminic acid, C.I. 75470

Natural Red 4. For staining mucines and glycogen.

5 g	CARA-P-5
10 g	CARA-P-10
25 g	CARA-P-25

Celestine Blue, C.I. 51050

Celestine blue B, Mordant blue 14. For staining nuclei dark blue, used as hematoxylin substitute.

10 g	CB-P-10
25 g	CB-P-25

Chromotrope 2R, C.I. 16570

Acid Red 29, Mordant Blue 80. For counterstaining cytoplasm in histopathology used as eosin substitute in HE staining and in trichrome stains.

25 g	C2R-P-25
------	----------

Congo Red, C.I. 22120

Direct Red 28, Kongorot, BSC certified stain. For staining of amyloids according to Highman and Puchtler and for elastic fiber staining.

25 g	CR-P-25
------	---------

Cresyl Violet Acetate Certified

Cresyl Violet, Oxazine 9, Cresyl Echt Violet. For use in microbiology and histology.

25 g	CVA-P-25
------	----------

Crystal Violet, C.I. 42555

Gentian Violet, Methyl Violet 10B, BSC certified stain. For bacterial staining according to Gram.

25 g	CRV-P-25
50 g	CRV-P-50

Eosin B, C.I. 45400

Eosin B, Eosin Bluish, Imperial Red, Acid Red 91, BSC certified. For staining according to Romanowsky. Used for staining red blood cells, connective tissues and as counterstain for the Gram stain.

25 g	EB-P-25
------	---------

Eosin Y, C.I. 45380

Eosin Yellowish, Acid Red 87, Eosin WS, BSC certified stain. Counterstain for hematoxylin and for preparing EA Pap stains and polychromatic hematological stains.

25 g	EOY-P-25
100 g	EOY-P-100
500 g	EOY-P-500

Eriochrome Cyanine R, C.I. 43820

Mordant Blue 3, Chromoxane Cyanine R. For use in cytology and histology.

25 g	ECR-P-25
------	----------

Fast Green F.C.F., C.I. 42053

Food Green 3, Solid Green F.C.F, 1724 Green, BSC certified stain. For trichrome staining according to Gomori and Masson.

25 g	FG-P-25
------	---------

Fuchsin Acid, C.I. 42685

Acid fuchsine, Acid Violet 19, BSC certified stain. For staining connective tissue according to Van Gieson and Mallory.

25 g	FA-P-25
------	---------

Fuchsin Basic, C.I. 42510

Basic fuchsine, Basic Violet 14, BSC certified stain. For staining mucins, Negri bodies and acid-fast bacteria.

25 g	FB-P-25
100 g	FB-P-100

Fuchsin Basic - Schiff, C.I. 42500

Pararosaniline HCl, Basic Red 9, BSC certified stain. Used for P.A.S. (Periodic Acid Schiff) staining, DNA staining according to Feulgen and acid-fast bacteria staining.

25 g	FBS-P-25
50 g	FBS-P-50

Gentian Violet, C.I. 42555

Crystal Violet, Methyl Violet 10B, BSC certified stain. For bacterial staining according to Gram.

25 g	GV-P-25
100 g	GV-P-100

Giemsa

BSC certified stain. Romanowsky dye for staining according to Pappenheim. Polychromatic dye used for identification of chromosomal damage and as a differentiating stain.

25 g	G-P-25
50 g	G-P-50