

Technical Data

Diphtheria Virulence Supplement (Part A & Part B)

FD073

A selective supplement recommended for the isolation and presumptive identification of Corynebacterium diphtheriae.

Composition

Per vial sufficient for 1000 ml medium

Ingredients Concentration

Part A

Horse serum 100ml

Part B

Potassium tellurite 1ml

Directions:

Warm up the refrigerated contents of Part B vial and aseptically add 29 ml sterile distilled water. Mix thoroughly. Aseptically add warmed up (to 50°C) contents of Part A and B vials to sterile, molten, cooled (45-50°C) Tinsdale Agar Base M314 / Tinsdale HiVegTM Agar Base MV314 as required.

For 10 ml of M314: 1.0 ml of Part A and 0.3 ml of Part B, is recommended.

Mix well and pour into sterile petri plates.

Storage and Shelf Life

Store Part A at -20°C & Part B at 2-8°C. Use before the expiry date on the label.

Revision: 1 / 2012

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia[™] publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia[™] Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.