



Technical Data

HiCrome™ Vibrio Agar

M1682

Intended use

Recommended for the isolation, and selective chromogenic differentiation of *Vibrio* species from seafood and clinical samples.

Composition**

Ingredients	Gms / Litre
Peptone	10.000
Sodium chloride	25.000
Sodium thiosulphate	5.000
Sodium citrate	6.000
Sodium cholate	1.000
Chromogenic mixture	5.500
Agar	15.000
Final pH (at 25°C)	8.5±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 67.5 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 45-50°C. Mix well before pouring into sterile Petri plates.

Principle And Interpretation

Vibrio's have played a significant role in human history. Outbreaks of cholera, caused by *Vibrio cholerae*, can be traced back in time to early recorded descriptions of enteric infections. The *Vibrio*'s have also received the attention of marine microbiologists who observed that the readily cultured bacterial population in near-shore waters and those associated with fish and shell fish were predominantly *Vibrio* species (7). *Vibrio* species are mainly responsible for causing cholera and food poisoning in humans. *Vibrio cholerae* causes cholera due to the intake of contaminated food such as raw oysters. *Vibrio parahaemolyticus* is a major cause of food borne infections, causing food poisoning (1). Since *Vibrio* species naturally occur in sea water, worth special mention is their need for sodium chloride, although some species can grow with minimum sodium chloride concentration (7). The widely used media for *Vibrio* isolation are TCBS Agar and Alkaline Peptone Water (2). However accompanying sucrose-fermenting bacteria pose a problem in the identification of *Vibrio* species on TCBS Agar. On HiCrome™ Vibrio Agar, the colour development by *Vibrio* species is not affected by the presence of colonies of other bacteria. This is because, the amount of colour developed depends on the reaction of the bacterial beta-galactosidase with the substrate contained in the media (4).

Peptone provides carbonaceous, nitrogenous and essential nutrients to the organisms. High concentration of sodium chloride in addition to maintaining the osmotic equilibrium also has an inhibitory action on the accompanying microflora. Sodium thiosulphate, sodium citrate and sodium cholate are used in the formulation because they can inhibit the growth of gram positive and some gram negative bacteria, but not members of *Enterobacteriaceae*. The proprietary chromogenic mixture incorporated in the medium helps in the chromogenic differentiation of *Vibrio cholerae* and *Vibrio parahaemolyticus*. The high (alkaline) pH of the medium helps in selective isolation of *Vibrio* species.

Type of specimen

Clinical samples- faeces; Food samples.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (3,5). For food samples, follow appropriate techniques for sample collection and processing as per guidelines (6). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic Use. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Being highly selective, some species may show poor growth due to nutritional variations.
2. Slight colour variation may be observed depending upon strains.
3. Further biochemical tests must be carried out for confirmation.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to light tan homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 6.75% w/v aqueous solution at 25°C. pH : 8.5±0.2

pH

8.30-8.70

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Vibrio cholerae</i> ATCC 15748	50-100	good-luxuriant	≥50%	purple
<i>Vibrio parahaemolyticus</i> ATCC 17802 (00037*)	50-100	good-luxuriant	≥50%	bluish green
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	≥10 ⁴	inhibited	0%	
<i>Escherichia coli</i> ATCC 25922 (00013*)	≥10 ⁴	inhibited	0%	
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	≥10 ⁴	inhibited	0%	

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (3,5).

Reference

1. Alcamo. E.I, 2001. Fundamentals of Microbiology, 6th ed, Jones and Bartlett Publishers, Inc. pg 254, 244.
2. Clesceri, Greenberg and Eaton (ed.). 1998. Standard Method for the examination of Water and Waste water, 20th ed. American Public Health Association, Washington, D. C.
3. Isenberg, H. Clinical Microbiology Procedures Handbook 2nd Edition.
4. Kudo. H. Y et al, 2001. Improved Method for Detection of *Vibrio parahaemolyticus* in Seafood. ASM. Vol 67, No.12, pg 5819-5823.
5. Murray P. R., Baron J. H., Pfaller M. A., Tenover J. C. and Tenover F. C., (Eds.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
6. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of, Foods, 5th Ed., American Public Health Association, Washington, D.C.
7. Thompson et al (ed.). 2006. The Biology of Vibrios, ASM Press, chapter 1, pg 3.


Revision : 03/2019



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India



CE Partner 4U, Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.