



Thayer Martin Medium Base

M413

Thayer Martin Medium Base used for selective isolation of Gonococci from pathological specimens.

Composition**

Ingredients	Gms / Litre
Peptone, special	23.000
Starch	1.000
Sodium chloride	5.000
Agar	13.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 21.0 grams in 450 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45°C. Aseptically add 50 ml of sterile lysed blood and rehydrated contents of one vial of Vitamino Growth Supplement (FD025) and V.C.N Supplement (FD023) or V.C.N.T Supplement (FD024). If desired GC Supplement with Antibiotics (FD021) can be used as a single supplement. Mix well before pouring into sterile Petri plates. If Hemoglobin (FD022) is used suspend 21.0 grams of Thayer Martin Medium Base in 250 ml distilled water. Heat to boiling to dissolve the medium completely. Prepare 250 ml of 2% hemoglobin solution. Sterilize separately by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45°C. Mix both and add the supplements as above.

Principle And Interpretation

Carpenter and Morton reported an improved medium to isolate Gonococci in 24 hours (1). Later on the efficiency of GC medium supplemented with haemoglobin and yeast concentrate was demonstrated for isolating gonococci (2). Subsequently Thayer and Martin Medium was developed for the primary isolation of *Neisseria gonorrhoeae* and *Neisseria meningitidis* from specimens containing mixed flora collected from throat, vagina, rectum and urethra (3, 4). Thayer and Martin (4) used Vancomycin, Colistin and Nystatin. Martin and Lester (5) used an additional antibiotic Trimethoprim to make the medium selective.

Special peptone provides nutrients to the organisms while starch neutralizes the toxic fatty acids if present in the agar. Haemoglobin provides the X factor whereas the V factor (N.A.D.) is provided by the added supplement. Supplement (FD025) also supplies vitamins, amino acids, coenzymes etc. which enhances the growth of pathogenic *Neisseria*. Vancomycin and colistin inhibits gram-positive and gram-negative bacteria respectively (6). Nystatin inhibits fungi. This medium may inhibit *Haemophilus* species. Some strains of *Capnocytophaga* species may grow on this medium when inoculated with oropharyngeal specimens

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.3% Agar gel.

Colour and Clarity of prepared medium

Basal Medium : Yellow coloured clear to slightly opalescent gel. After addition of haemoglobin or sterile lysed blood and supplements: chocolate coloured opaque gel forms in Petri plates.

Reaction

Reaction of 4.2% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

M413: Cultural characteristics observed with added sterile lysed blood/Haemoglobin solution (FD022), Vitamino Growth Supplement (FD025) and V.C.N. Supplement (FD023)/V.C.N.T. Supplement (FD024) after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Escherichia coli</i> ATCC 25922	$\geq 10^3$	inhibited	0%	
<i>Neisseria gonorrhoeae</i> ATCC 19424	50-100	good-luxuriant	$\geq 50\%$	small, grayish-white to colourless, mucoid
<i>Neisseria meningitidis</i> ATCC 13090	50-100	good-luxuriant	$\geq 50\%$	medium to large, blue-gray, mucoid
<i>Proteus mirabilis</i> ATCC 25933	$\geq 10^3$	inhibited	0%	

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. Carpenter and Morton, 1947, Proc. N.Y. State Assoc. Public Hlth. Labs., 27:58.
2. Carpenter et al, 1949, Am. J. Syphil. Gonorrh. Vener. Dis., 33:164.
3. Martin, Billings, Hackney and Thayer, 1967, Public Hlth. Rep., 82:361.
4. Thayer J. and Martin J.E. Jr., 1966, Public Health Rep., 81:559.
5. Martin J.E. Jr. and Lester A., 1971, HSMHA Hlth. Service Rep., 86(1):30.
6. MacFaddin J., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.