




MICROSCOPY AND
LABORATORY DIAGNOSTICS

Our global presence

“ We manufacture products for use in histopathology, hematology, cytology, and microbiology with a wide national and international distribution network present in over 70 countries.

BioGnost offers only the best quality products at competitive prices and has one of the widest ranges of medical products in South-Eastern Europe. Our experience and flexibility towards the client makes us a sought-after business partner, which has been recognized on the global market.



Our entrepreneurship started in 1990 with the founding of the family company. Accelerated business growth began in 1998 with production of blood typing serums followed by the Croatian Ministry of Health granting BioGnost the License for the manufacture of medical products. We are proud of one of our biggest successes - relocating to our new commercial and manufacturing building in 2012. The site enabled an enhanced work environment, including a carefully devised warehouse structure, segmented production facilities with increased manufacturing capacity, and additional R&D capabilities.

“ We continue to build on the strong momentum from last year to create high-quality products and provide excellent customer service.


Strategic priorities

Our mission is to create high-quality products and provide excellent services, as well as introduce new or expanded product lines and sales ranges in accordance with market dynamics and demands.

Product and service excellence makes BioGnost a desirable regional and international business partner. The key to success is the ever-expanding network of business partnerships, as well as recognizable social responsibility. In the coming years we plan to further our market expansion, additionally invest in research and development, and to become a competitive manufacturer in the fields of microbiological diagnostics and histopathology.

ISO 9001, ISO 13485, ISO 14001

BioGnost's entire business is in accordance with international ISO 9001, ISO 13485 and ISO 14001 standards. The company implemented ISO standards more than 15 years ago, and is the first Croatian company to implement the ISO 13485 quality system for medical devices. Having fulfilled the requirements of these standards, the company is completely in accordance with national legal regulation and the European Union's Medical Device Regulation (EU 2017/745).



CE, IVDR

BioGnost's products bear the CE mark, a certificate that confirms that the products are in accordance with the European Union's standards. They are declared as *in vitro* diagnostic medical devices (IVD) classified as "Class A" products (according to the Regulation (EU) 2017/746). No control of a Notified Body is required for this class; entire company business is in accordance with applicable regulations and standards.

OEM manufacture

Besides products from our own sales range, we also offer OEM manufacture for your company. It includes a wide range of services, such as label design, product packaging, instructions for use and certificates of analysis, and products manufactured according to your specifications.

Contents



FIXATION AND PROCESSING 6

Marking systems 6

Formalin fixatives 8

Other fixatives 10

Decalcifying solutions 11

Dehydrating and rehydrating agents 12

Clearing and deparaffinization agents 13

Infiltration, embedding and demolding media 14

MOUNTING AND MICROSCOPY 16

Mounting media 16

Immersion oils 20

Cleaning solutions for microscope lens and slides 21

STANDARD HISTOLOGY STAINING REAGENTS 22

Hematoxylin reagents 22

Differentiation and bluing reagents 24

Eosin reagents 25

CYTOLOGY AND HEMATOLOGY STAINING REAGENTS 26

Papanicolaou (Pap) reagents 26

Rapid and standard hematology reagents and kit 28

LEUKOGNOST Cytochemical staining kits and ancillary reagents 32

BACTERIOLOGY AND MYCOLOGY STAINING REAGENTS 34

BioGram kits and reagents 34

TB-Stain kits and reagents 36

Solutions for staining and analysis of fungi 39

SPECIAL STAINING KITS FOR HISTOLOGY AND CYTOLOGY 40

ADDITIONAL STAINING REAGENTS AND SOLUTIONS 51

POWDER DYES AND STAINS 58

ACCESSORIES 62

Legend:

● DG cargo in transport ○ DG cargo only in air transport ● temperature controlled cargo (transport regime 2-8°C)

Fixation and processing

The initial part of clinical sample processing consists of products necessary for successful processing and obtaining optimal final results. That includes fixatives that preserve the tissue's structure, alcohols, paraffins that enable section cutting and finally, immersion oils for microscopy and mounting media for long-term storage of slides.

Marking systems

Marking systems consist of dyes for permanent tissue marking and markers used for marking glass slides, cassettes and transport containers. Tissue marking dyes are used for marking and orientation of tissue samples intended for further histopathology processing.



Tissue marking dyes, 5 color set

TMD Tissue marking dyes set 5x50 mL contains 5 dyes for tissue marking: green, yellow, black, red and blue. The set includes wooden sticks and rack.

5x50 mL 3408-SET

Tissue marking dyes, 7 color set

TMD Tissue marking dyes set 7x50 mL contains 7 dyes for tissue marking: green, yellow, black, red, blue, orange and lilac. The set includes wooden sticks and rack.

7x50 mL 3408-SET-7

Tissue marking dyes

For marking and orientation of tissue samples.

| | | |
|----------------------------|--------|------------|
| Tissue marking dye, green | 50 mL | 3408-1-50 |
| | 100 mL | 3408-1-100 |
| | 250 mL | 3408-1-250 |
| Tissue marking dye, yellow | 50 mL | 3408-2-50 |
| | 100 mL | 3408-2-100 |
| | 250 mL | 3408-2-250 |
| Tissue marking dye, black | 50 mL | 3408-3-50 |
| | 100 mL | 3408-3-100 |
| | 250 mL | 3408-3-250 |
| Tissue marking dye, red | 50 mL | 3408-4-50 |
| | 100 mL | 3408-4-100 |
| | 250 mL | 3408-4-250 |
| Tissue marking dye, blue | 50 mL | 3408-5-50 |
| | 100 mL | 3408-5-100 |
| | 250 mL | 3408-5-250 |
| Tissue marking dye, orange | 50 mL | 3408-6-50 |
| | 100 mL | 3408-6-100 |
| | 250 mL | 3408-6-250 |
| Tissue marking dye, lilac | 50 mL | 3408-7-50 |
| | 100 mL | 3408-7-100 |
| | 250 mL | 3408-7-250 |

Liquid Blocker (PAP pen)

Liquid Blocker contains hydrophobic liquid as a water repellent barrier around specimens or sections. Useful in immunohistochemistry that uses small amounts of antiserum. Multiple specimens can be separated by drawn circles or lines applied to the same slide.

1 pc ML-1

Duoline marker pen

Professional marker pen for permanent labelling of cassettes, slides and transport containers. With fine and thick marker pen tip.

1 pc MD-1





Formalin fixatives

Fixation is one of the most important steps in tissue processing, preventing tissue degradation and decomposition. Preserving tissue properties is important for detection of morphological and pathological changes in tissue samples.

4% NB Formaldehyde

10% neutral buffered and stabilized formalin solution, pH 7.0.

| | |
|----------|----------|
| 1000 mL | FNB4-1L |
| 5000 mL | FNB4-5L |
| 10000 mL | FNB4-10L |
| 20000 mL | FNB4-20L |



4% Formaldehyde

Stabilized 10% formalin solution.

| | |
|----------|--------|
| 1000 mL | F4-1L |
| 5000 mL | F4-5L |
| 10000 mL | F4-10L |
| 20000 mL | F4-20L |



10% NB Formaldehyde

25% neutral buffered and stabilized formalin solution, pH 7.0.

| | |
|----------|-----------|
| 1000 mL | FNB10-1L |
| 5000 mL | FNB10-5L |
| 10000 mL | FNB10-10L |
| 20000 mL | FNB10-20L |



ImmunoForm

Stabilized 10% formalin solution produced from paraformaldehyde without methyl alcohol. For use in histology and immunohistochemistry.

| | |
|----------|---------|
| 1000 mL | IMF-1L |
| 5000 mL | IMF-5L |
| 10000 mL | IMF-10L |



10% Formaldehyde

Stabilized 25% formalin solution.

| | |
|----------|---------|
| 1000 mL | F10-1L |
| 5000 mL | F10-5L |
| 10000 mL | F10-10L |
| 20000 mL | F10-20L |



Formaldehyde 37% for use in histology

100% formalin solution stable at low temperatures.

| | |
|----------|----------|
| 1000 mL | F37H-1L |
| 5000 mL | F37H-5L |
| 10000 mL | F37H-10L |



NB Formaldehyde concentrated 5x

Concentrated neutral buffered formalin for preparation of 10% buffered formalin (synonym: 4% buffered formaldehyde) solution, pH 7.0.

| | | |
|--|---------|---------|
| For preparation of 5 L of 4% buffered formaldehyde | 1000 mL | FK5X-1L |
|--|---------|---------|

| | | |
|---|---------|---------|
| For preparation of 25 L of 4% buffered formaldehyde | 5000 mL | FK5X-5L |
|---|---------|---------|

| | | |
|---|----------|----------|
| For preparation of 50 L of 4% buffered formaldehyde | 10000 mL | FK5X-10L |
|---|----------|----------|



DeHematin

Modified Kardasewitsch solution for depigmentation of tissues fixed in non-buffered formalin (after acid hematin occurs).

| | |
|---------|----------|
| 1000 mL | KO-OT-1L |
| 5000 mL | KO-OT-5L |



Other fixatives

BioFix

Methyl alcohol-based fixative for cryostat sections.

| | |
|---------|--------|
| 500 mL | BF-500 |
| 1000 mL | BF-1L |



BioFix C

Modified Carnoy's fixative based on ethyl alcohol, chloroform and acetic acid.

| | |
|---------|---------|
| 500 mL | BFC-500 |
| 1000 mL | BFC-1L |
| 5000 mL | BFC-5L |



BioFix GL

Glyoxal-based fixative for use in histology.

| | |
|---------|----------|
| 500 mL | BFGL-500 |
| 1000 mL | BFGL-1L |
| 5000 mL | BFGL-5L |

BioFix HD

Modified Hartmann/Davidson's fixative that enables rapid fixation and better visualization of lymph nodes. The results provide clearer nuclear details and minimize formalin pigment occurrence.

| | |
|---------|----------|
| 500 mL | BFHD-500 |
| 1000 mL | BFHD-1L |
| 5000 mL | BFHD-5L |



BioForm S

Modified Schaeffer's buffered fixative solution of formaldehyde in methanol.

| | |
|---------|------------|
| 1000 mL | BIOS-OT-1L |
| 5000 mL | BIOS-OT-5L |





Other fixatives

Bouin's solution

Aqueous solution of picric acid, formaldehyde and acetic acid for tissue fixation and better visualization of lymph nodes.

| | |
|----------|------------|
| 100 mL | BOU-OT-100 |
| 1000 mL | BOU-OT-1L |
| 5000 mL | BOU-OT-5L |
| 10000 mL | BOU-OT-10L |



CitoFix

Alcoholic solution for fixation and transport of cytological samples.

| | |
|---------|-------|
| 1000 mL | CX-1L |
|---------|-------|



CitoSpray

Solution for fixation of cytological smears in a spray bottle. Contains polyglycol which forms protective film when applied on cytological smear.

| | |
|--------|--------|
| 100 mL | CS-100 |
|--------|--------|



CitoSpray solution

Solution for fixation of cytological smears in a bottle, for refill of CitoSpray bottle.

| | |
|---------|-------|
| 1000 mL | CS-1L |
|---------|-------|



Saccomanno's fixative

Solution for fixation and transport of cytological samples.

| | |
|---------|-----------|
| 500 mL | SF-OT-500 |
| 1000 mL | SF-OT-1L |
| 5000 mL | SF-OT-5L |



Decalcifying solutions

Decalcifiers most often contain inorganic or organic acids enabling rapid decalcification (i.e. calcium removal). Milder types of decalcifiers are formulated from chelating agents that are used for decalcification of more sensitive types of tissue.

OsteoFast 1

Light blue fixative and decalcifying solution for bone and hard tissue in histology, based on formaldehyde and hydrochloric acid.

1000 mL OF1-OT-1L
2500 mL OF1-OT-2.5L



OsteoFast 2

Light yellow decalcifying solution for bone and hard tissue in histology, based on formic acid.

1000 mL OF2-OT-1L
2500 mL OF2-OT-2.5L



OsteoSens

Decalcifying solution for sensitive calcified tissues in histology, contains EDTA.

1000 mL OS-OT-1L
2500 mL OS-OT-2.5L





Dehydrating and rehydrating agents

Alcoholic solutions and absolute alcohol are ideal for histocytology and cytology processing, storing and transporting samples. They are also used for dehydration and rehydration of tested samples.

Histanol 50

Denaturated 50% ethyl alcohol.

| | |
|----------|---------|
| 1000 mL | H50-1L |
| 5000 mL | H50-5L |
| 10000 mL | H50-10L |



Histanol 70

Denaturated 70% ethyl alcohol.

| | |
|----------|---------|
| 1000 mL | H70-1L |
| 5000 mL | H70-5L |
| 10000 mL | H70-10L |



Histanol 80

Denaturated 80% ethyl alcohol.

| | |
|----------|---------|
| 1000 mL | H80-1L |
| 5000 mL | H80-5L |
| 10000 mL | H80-10L |



Histanol 95

Denaturated 95% ethyl alcohol.

| | |
|----------|---------|
| 1000 mL | H95-1L |
| 5000 mL | H95-5L |
| 10000 mL | H95-10L |



Histanol 96

Denaturated 96% ethyl alcohol.

| | |
|----------|---------|
| 1000 mL | H96-1L |
| 5000 mL | H96-5L |
| 10000 mL | H96-10L |



Histanol 100

Denaturated 100% ethyl alcohol.

| | |
|----------|----------|
| 1000 mL | H100-1L |
| 5000 mL | H100-5L |
| 10000 mL | H100-10L |



Histanol EP 95

95% mixture of denaturated ethyl alcohol and isopropyl alcohol.

| | |
|----------|-----------|
| 1000 mL | HEP95-1L |
| 5000 mL | HEP95-5L |
| 10000 mL | HEP95-10L |



Histanol EP 100

100% mixture of denaturated ethyl alcohol and isopropyl alcohol.

| | |
|----------|------------|
| 1000 mL | HEP100-1L |
| 5000 mL | HEP100-5L |
| 10000 mL | HEP100-10L |



Histanol EP 9010 NEW

Alcoholic solvent substituting standard ethanol series in processing, dewaxing and dehydrating.

| | |
|----------|-----------|
| 1000 mL | HEP91-1L |
| 5000 mL | HEP91-5L |
| 10000 mL | HEP91-10L |



Histanol IP

Isopropyl alcohol for use in microscopy.

| | |
|----------|---------|
| 1000 mL | HIP-1L |
| 5000 mL | HIP-5L |
| 10000 mL | HIP-10L |



Histanol M

Methyl alcohol for use in microscopy.

| | |
|----------|--------|
| 1000 mL | HM-1L |
| 5000 mL | HM-5L |
| 10000 mL | HM-10L |



Histanol G

Glycerol for use in microscopy.

| | |
|----------|--------|
| 1000 mL | HG-1L |
| 5000 mL | HG-5L |
| 10000 mL | HG-10L |



Clearing and deparaffinization agents

Clearing agents are completely miscible with alcohol and paraffin, making them an ideal intermedium in histology and cytology processing. They are used multiple times during tissue processing and section staining; before embedding the tissue in paraffin, for deparaffinization before staining and before covering the stained section using medium for permanent preservation.

BioClear

Xylene-based medium for use in histology and cytology.

| | |
|----------|---------|
| 1000 mL | BC-1L |
| 2500 mL | BC-2.5L |
| 5000 mL | BC-5L |
| 10000 mL | BC-10L |

BioClear New

Xylene substitute, aliphatic hydrocarbon-based. Essentially no odor. For use in histology and cytology.

| | |
|----------|----------|
| 1000 mL | BCN-1L |
| 2500 mL | BCN-2.5L |
| 5000 mL | BCN-5L |
| 10000 mL | BCN-10L |





Infiltration, embedding and demolding media

Paraffin wax is the most commonly used sample for infiltration, embedding and forming tissue blocks. It provides support to the tissue during microtome cutting.

BioWax 52/54

Blend of paraffin waxes with low melting point at 52/54°C for processing wide range of different samples. Lower melting point makes it suitable for working with delicate tissues without deforming or damaging and allows good preservation of tissue morphology during processing.

| | |
|-------|------------|
| 1 kg | BW52/54-1 |
| 2 kg | BW52/54-2 |
| 10 kg | BW52/54-10 |

BioWax 56/58

Blend of paraffin waxes with melting point at 56/58°C for optimal infiltration and routine work, enables easier formation of ribbons, virtually no cut resistance and better overall results.

| | |
|-------|------------|
| 1 kg | BW56/58-1 |
| 2 kg | BW56/58-2 |
| 10 kg | BW56/58-10 |

BioWax Blue

Blend of blue paraffin waxes with melting point at 56/58°C and added plastic polymers for optimal infiltration. Especially suitable for small tissue samples. Blue color provides better contrast enabling easier visualization during processing. It is also recommended for use in immunoperoxidase methods.

| | |
|-------|--------|
| 1 kg | BWB-1 |
| 2 kg | BWB-2 |
| 10 kg | BWB-10 |

BioWax Plus 56/58

An optimal blend of paraffin wax and plastic polymers with melting point at 56/58°C. Its properties include low shrinking rate during cooling down and excellent section creation without deformations. Provides excellent infiltration into tissues and reduces the tendency of tissue to crack.

| | |
|-------|-----------|
| 1 kg | BWPLUS-1 |
| 2 kg | BWPLUS-2 |
| 10 kg | BWPLUS-10 |

Micro Clean

Solution in spray bottle for rapid and efficient paraffin residue removal from work surfaces and instruments.

| | |
|--------|-----------|
| 100 mL | MC-OT-100 |
|--------|-----------|



CryoFix Gel

Embedding matrix for cryosectioning of tissue samples known as OCT medium (optimal cutting temperature). Efficient at temperatures of -10°C and below. Available in 5 different colors.

| | | |
|--------------------|--------|---------|
| CryoFix Gel | 100 mL | CF-100 |
| CryoFix Gel pink | 100 mL | CFR-100 |
| CryoFix Gel blue | 100 mL | CFP-100 |
| CryoFix Gel green | 100 mL | CFZ-100 |
| CryoFix Gel yellow | 100 mL | CFY-100 |



Cryo Spray

Medium in spray bottle for freezing histological samples.

200 mL 08-SPRAY-2



DeMold Spray

Solution in spray bottle for easy removal of paraffin tissue blocks from the mold.

100 mL DS-100



DeMold solution

Solution for refilling DeMold Spray bottle.

500 mL DS-500
1000 mL DS-1L



Tissue Soft

Solution for softening the tissue embedded in paraffin block before microtome sectioning.

500 mL TS-500
1000 mL TS-1L



Mounting and microscopy

Protection and preservation of stained samples generally require mounting with a suitable medium. For further examination of the sample, immersion oils are essential in providing a high resolution image during microscopical analysis at high magnification. Special cleaning solutions are used for removing oil residues from microscope lens and slides.

Mounting media

Ten different covering and mounting media for permanent preservation of microscope slides

In order to prepare permanent sections, stained samples must be covered with a suitable mounting medium. One of the most important criteria is the refractive index similar to that of glass. Other criteria include a suitable solvent, polymer and optimal viscosity. After the sample is covered and a cover glass is mounted, the section is protected from external factors and remains preserved for a prolonged period of time. BioMount media do not crack or turn yellow during sample storage.



BioGnost's synthetic BioMount media properties

| PRODUCT NAME | SOLVENT | POLYMER | VISCOSITY @ 20°C (cSt, mPa) | REFRACTIVE INDEX @ 20°C (n _p) | USAGE IN AUTOMATED COVER SLIP MACHINES |
|-------------------|----------------------------|---------|--------------------------------|--|---|
| BIOMOUNT DPX | xylene (BioClear) | S | 850-950 | 1.5210-1.5230 | - |
| BIOMOUNT DPX HIGH | xylene (BioClear) | S | 1900-2100 | 1.5235-1.5255 | - |
| BIOMOUNT DPX LOW | xylene (BioClear) | S | 140-160 | 1.5115-1.5135 | ✓ |
| BIOMOUNT DPX NEW* | xylene (BioClear) | S | 200-300 | 1.5150-1.5180 | ✓ |
| BIOMOUNT NEW | isoparaffin (BioClear New) | M | 950-1050 | 1.4385-1.4395 | - |
| BIOMOUNT NEW LOW | isoparaffin (BioClear New) | M | 130-160 | 1.4345-1.4365 | ✓ |
| BIOMOUNT | xylene (BioClear) | M | 450-650 | 1.4920-1.4930 | ✓ |
| BIOMOUNT HIGH | xylene (BioClear) | M | 5000-5500 | 1.4900-1.4920 | - |
| BIOMOUNT M | toluene | M | 140-160 | 1.4900-1.4920 | ✓ |
| BIOMOUNT C | xylene (BioClear) | M | 5000-5500 | 1.4910-1.4925 | - |

Legend: M – methacrylate, S – styrene (polystyrene)
 * does not contain harmful dibutyl phthalate plasticizer
 ** contact us to find out how to decrease viscosity of a medium

Xylene and styrene-based media for permanent mounting

BioMount DPX

Moderate viscosity medium for permanent covering of microscope samples and mounting cover glass.

100 mL BM-100
 250 mL BM-250
 500 mL BM-500



BioMount DPX High

High viscosity medium for permanent covering of microscope samples and mounting cover glass.

100 mL BMH-100
 250 mL BMH-250
 500 mL BMH-500



BioMount DPX Low

Low viscosity medium for permanent covering of microscope samples and mounting cover glass. Can be used in automatic cover slip machines.

100 mL BML-100
 250 mL BML-250
 500 mL BML-500



BioMount DPX New

Low viscosity medium for permanent covering of microscope samples and mounting cover glass. Medium contains more environmentally friendly plasticizer instead of harmful dibutyl phthalate. Can be used in automatic cover slip machines.

100 mL BMLE-100
 250 mL BMLE-250
 500 mL BMLE-500





Methacrylate and xylene substitute-based media for permanent mounting

BioMount New

Moderate viscosity medium for permanent covering of microscope samples deparaffinized using xylene substitute-based medium, as well as xylene-based medium.

| | |
|--------|---------|
| 100 mL | BMN-100 |
| 250 mL | BMN-250 |
| 500 mL | BMN-500 |

BioMount New Low

Very low viscosity medium for permanent covering of microscope samples deparaffinized using xylene substitute, as well as xylene.

| | |
|--------|----------|
| 100 mL | BMNL-100 |
| 250 mL | BMNL-250 |
| 500 mL | BMNL-500 |



Methacrylate-based permanent mounting media

BioMount

Low viscosity xylene-based medium for permanent covering of microscope slides and mounting cover glass.

| | |
|--------|---------|
| 100 mL | BMT-100 |
| 250 mL | BMT-250 |
| 500 mL | BMT-500 |



BioMount M

Very low viscosity toluene-based medium for permanent covering of microscope samples and mounting cover glass.

| | |
|--------|---------|
| 100 mL | BMM-100 |
| 250 mL | BMM-250 |
| 500 mL | BMM-500 |



BioMount High

Very high viscosity xylene-based medium for permanent covering of microscope samples and mounting cover glass.

| | |
|--------|----------|
| 100 mL | BMTH-100 |
| 250 mL | BMTH-250 |
| 500 mL | BMTH-500 |



Additional permanent mounting media

BioMount C

Very high viscosity medium for permanent covering of microscope slides and mounting cover glass.

| | |
|--------|---------|
| 100 mL | BMC-100 |
| 250 mL | BMC-250 |
| 500 mL | BMC-500 |

* excellent synthetic substitute for natural covering media, such as Canada Balsam



Water-based mounting media

| PRODUCT NAME | SOLVENT | POLYMER | VISCOSITY @ 20°C (cSt, mPa) | REFRACTIVE INDEX @ 20°C (n _p) | USAGE IN AUTOMATED COVER SLIP MACHINES |
|---------------|---------|---------|-----------------------------|---|--|
| BIOMOUNT AQUA | water | PVP | 700-1000 | 1.4145-1.4205 | - |



BioMount Aqua

Water-based medium for covering microscope slides and mounting cover slips. With application in hematology, immunohistochemistry and immunocytochemistry with Fast-Red or AEC chromogens that require an aqueous mounting medium, but also with DAB chromogen.

| | |
|--------|---------|
| 30 mL | BMA-30 |
| 100 mL | BMA-100 |
| 250 mL | BMA-250 |
| 500 mL | BMA-500 |



Immersion oils

Seven different types of immersion oils for microscopical analysis

Immersion oils are essential for routine laboratory work, especially in hematology, cytology and microbiology. They are commonly used for high magnifications in order to secure high resolution and a clear image. There are several types of immersion oils that differ according to viscosity and certain properties required by different microscopy techniques. They all have a common refractive index of 1.5150, equal to the refractive index of glass.

BioGnost's Immersion oils properties

| PRODUCT NAME | VISCOSITY @ 23°C (cSt, mPa) | COMBINING WITH OTHER TYPES OF OIL | TYPE OF MICROSCOPY |
|-------------------------------------|--------------------------------|--|---------------------------|
| IMMERSION OIL | 800-1300 | Immersion oil type A Immersion oil type C | light |
| IMMERSION OIL TYPE A | 100-300 | Immersion oil Immersion oil type C | light |
| IMMERSION OIL TYPE C | 1500-2200 | Immersion oil Immersion oil type A | light |
| IMMERSION OIL TYPE FF | 170 | - | fluorescent |
| IMMERSION OIL TYPE 37* | 1250** | - | light, for use at 37°C |
| IMMERSION OIL, TROPICAL GRADE*** | 1000-1500**** | - | light, for use at 27-29°C |

* refractive index n_D at 37°C = 1.5150 ** viscosity at 37°C *** refractive index n_D at tropical temperatures = 1.515 **** viscosity at 28°C

Immersion oil

Immersion oil for microscopy analysis with optimal viscosity of 800-1300 cSt.

| | |
|----------|---------|
| 10x10 mL | IU-1010 |
| 30 mL | IU-30 |
| 100 mL | IU-100 |
| 250 mL | IU-250 |
| 500 mL | IU-500 |

Immersion oil type A

Immersion oil for microscopy analysis with low viscosity of 100-300 cSt.

| | |
|----------|----------|
| 10x10 mL | IUA-1010 |
| 30 mL | IUA-30 |
| 100 mL | IUA-100 |
| 250 mL | IUA-250 |
| 500 mL | IUA-500 |





Immersion oil type C

Immersion oil for microscopy analysis with high viscosity of 1500-2200 cSt.

100 mL IUC-100
250 mL IUC-250
500 mL IUC-500

* excellent synthetic substitute for natural immersion oils, such as Cedarwood oil



Immersion oil type FF

Immersion oil for sample analysis in fluorescent microscopy.

100 mL IUF-100
250 mL IUF-250
500 mL IUF-500

Immersion oil type 37

Immersion oil for microscopy analysis with optimal refractive index and viscosity of 1250 cSt at 37°C.

100 mL IU37-100
250 mL IU37-250
500 mL IU37-500



Immersion oil, tropical grade

Immersion oil for microscopy analysis in tropical climates with optimal refractive index at 27-29°C.

10x10 mL IUT-1010
30 mL IUT-30
100 mL IUT-100
250 mL IUT-250
500 mL IUT-500



Cleaning solutions for microscope lens and slides

Clean Lens, solution 1

Solution according to Carl Zeiss formulation for cleaning microscope lens. Based on n-hexane and 2-propanol.

100 mL CL1-OT-100



Clean Lens, solution 2

Solution for cleaning microscope lens and removing immersion oil from stained slide after using microscope. Based on diethyl ether and 2-propanol.

100 mL CL2-OT-100



Clean Lens, solution 3

Solution for cleaning microscope lens and removing immersion oil from stained slide after using microscope. Based on petroleum ether and 2-propanol.

100 mL CL3-OT-100





Standard histology staining reagents


Reagents for hematoxylin and eosin (HE) staining, the gold standard for demonstrating cellular and tissue structures, providing detailed anatomy and morphology of frozen and paraffin tissue sections, enabling clear visualization of nuclear, cytoplasmic and other cell structures.

Hematoxylin reagents

Nuclear stains for standard HE staining

Properties of BioGnost's Hematoxylin reagents

| PRODUCT NAME | CYTOLOGY | HISTOLOGY | IHC | PROGRESSIVE METHOD | REGRESSIVE METHOD |
|------------------------|----------|-----------|-----|--------------------|-------------------|
| HEMATOXYLIN H | - | ✓ | - | ✓ | ✓ |
| HEMATOXYLIN M | - | ✓ | ✓ | ✓ | - |
| HEMATOXYLIN ML | - | ✓ | ✓ | ✓ | ✓ |
| HEMATOXYLIN G1 | ✓ | ✓ | ✓ | ✓ | - |
| HEMATOXYLIN G2 | ✓ | ✓ | ✓ | ✓ | ✓ |
| HEMATOXYLIN G3 | ✓ | ✓ | - | ✓ | ✓ |
| HEMATOXYLIN CAR | - | ✓ | ✓ | ✓ | - |
| HEMATOXYLIN COL | - | ✓ | ✓ | ✓ | ✓ |
| HEMATOXYLIN HP, PAP 1A | ✓ | - | - | ✓ | ✓ |
| HEMATOXYLIN HP, PAP 1B | ✓ | - | - | ✓ | ✓ |
| HEMATOXYLIN W KIT | - | ✓ | ✓ | ✓ | acid resistant |
| HEMATOXYLIN P.T.A. | - | ✓ | - | ✓ | - |



Each of BioGnost's Hematoxylin is specially stabilized in order to maintain the reagent's reactivity for prolonged periods of time, as well as to maintain the quality of the reagent. Every staining provides a crystal clear and highly detailed image of the cell's nucleus, which in turn makes abnormal or pathological changes easier to detect.

Hematoxylin H

Modified hematoxylin according to Harris, most commonly used formulation in HE staining. Strong intensity reagent for progressive and regressive staining in histopathology.

| | |
|---------|--------------|
| 100 mL | HEMH-OT-100 |
| 500 mL | HEMH-OT-500 |
| 1000 mL | HEMH-OT-1L |
| 2500 mL | HEMH-OT-2.5L |

Hematoxylin M

Modified hematoxylin according to Mayer. Weaker intensity reagent for progressive staining in histopathology and counterstaining in immunohistochemistry.

| | |
|---------|--------------|
| 100 mL | HEMM-OT-100 |
| 500 mL | HEMM-OT-500 |
| 1000 mL | HEMM-OT-1L |
| 2500 mL | HEMM-OT-2.5L |

Hematoxylin ML

Modified hematoxylin according to Mayer-Lillie. Strong intensity new generation reagent for progressive and regressive staining in histopathology.

| | |
|---------|---------------|
| 100 mL | HEMML-OT-100 |
| 500 mL | HEMML-OT-500 |
| 1000 mL | HEMML-OT-1L |
| 2500 mL | HEMML-OT-2.5L |

Hematoxylin G1

Hematoxylin according to Gill, ideal for staining goblet cells. New generation reagent for delicate progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

| | |
|---------|---------------|
| 100 mL | HEMG1-OT-100 |
| 500 mL | HEMG1-OT-500 |
| 1000 mL | HEMG1-OT-1L |
| 2500 mL | HEMG1-OT-2.5L |

Hematoxylin G2

Hematoxylin according to Gill, contains double concentration of hematoxylin compared to Hematoxylin G1. Moderate intensity new generation reagent for progressive and regressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

| | |
|---------|---------------|
| 100 mL | HEMG2-OT-100 |
| 500 mL | HEMG2-OT-500 |
| 1000 mL | HEMG2-OT-1L |
| 2500 mL | HEMG2-OT-2.5L |

Hematoxylin G3

Modified hematoxylin contains triple concentration of hematoxylin compared to Hematoxylin G1. Strong intensity new generation reagent for progressive and regressive staining in histopathology and cytology.

| | |
|---------|---------------|
| 100 mL | HEMG3-OT-100 |
| 500 mL | HEMG3-OT-500 |
| 1000 mL | HEMG3-OT-1L |
| 2500 mL | HEMG3-OT-2.5L |





Hematoxylin reagents

Nuclear stains for standard HE staining

Hematoxylin W kit

Acid-resistant hematoxylin according to Weigert. Two-reagent kit that stains the nuclei blue-black, often a component of special staining kits for connective tissues.

| | |
|-----------|------------|
| 2x100 mL | HEMW-K-100 |
| 2x500 mL | HEMW-K-500 |
| 2x1000 mL | HEMW-K-1L |

Hematoxylin P.T.A.

Synonyms: Hematoxylin Phosphotungstic Acid, Mallory P.T.A.H. reagent. Simultaneously stains different types of tissue reddish-brown and blue. For differentiation between fibrin, smooth and cross-striated muscle fibers.

| | |
|---------|-------------|
| 500 mL | HPTA-OT-500 |
| 1000 mL | HPTA-OT-1L |

Hematoxylin CAR

Modified hematoxylin according to Carazzi for delicate nuclear staining without staining cytoplasmic components. Reagent for progressive staining of frozen sections in histopathology.

| | |
|---------|--------------|
| 100 mL | HEMCA-OT-100 |
| 500 mL | HEMCA-OT-500 |
| 1000 mL | HEMCA-OT-1L |

Hematoxylin COL

Modified hematoxylin according to Cole, contains iodine. Weaker intensity reagent for progressive and regressive staining in histopathology.

| | |
|---------|--------------|
| 100 mL | HEMCO-OT-100 |
| 500 mL | HEMCO-OT-500 |
| 1000 mL | HEMCO-OT-1L |

Hematoxylin Instant kit

Two-component kit that contains ready-to-use powders for preparation of hematoxylin reagent for progressive and regressive nuclear staining in histopathology and cytology.

| | |
|-------------|-----------|
| for 1000 mL | HEMI-K-1L |
| for 6000 mL | HEMI-K-6L |

Kit for rapid HE staining of frozen and paraffin-embedded sections

Ready-to-use eight-reagent kit (in 20 containers that can be used as staining jars) for rapid HE staining of frozen and paraffin tissue sections in histopathology. Contains xylene substitute as clearing agent and xylene substitute-based medium for permanent coverslipping.

| | |
|---------------|-------------|
| for 100 tests | HE-RTU-100T |
|---------------|-------------|

ErioGnost reagent

Acid fast reagent for nuclear staining, comparable to hematoxylin staining. Eco-friendly synthetic replacement for hematoxylin reagents.

| | |
|---------|------------|
| 100 mL | ERS-OT-100 |
| 500 mL | ERS-OT-500 |
| 1000 mL | ERS-OT-1L |

Differentiation and bluing reagents for histology and cytology

Differentiation of the nucleus enables clear visualization of nuclear structures in regressive procedure, while the bluing reagent turns the nucleus blue.

Acid alcohol

Reagent for enabling crisp visualization of nuclear chromatin detail and removing of excess background dye in histological samples after regressive staining with hematoxylin.

| | |
|---------|------------|
| 1000 mL | KA-OT-1L |
| 2500 mL | KA-OT-2.5L |

Scott's solution

Buffered reagent with potassium carbonate and magnesium sulphate for bluing of nuclear chromatin and nuclear membranes.

| | |
|---------|------------|
| 1000 mL | SC-OT-1L |
| 2500 mL | SC-OT-2.5L |

HCL Pap reagent

Aqueous reagent for enabling crisp visualization of nuclear chromatin detail and removing of excess background dye in cytological after regressive staining with hematoxylin.

| | |
|---------|----------------|
| 1000 mL | HCLPAP-OT-1L |
| 2500 mL | HCLPAP-OT-2.5L |

Bluing reagent

Blue colored buffered reagent with lithium carbonate for bluing of nuclear chromatin and nuclear membranes.

| | |
|---------|------------|
| 1000 mL | BR-OT-1L |
| 2500 mL | BR-OT-2.5L |

Eosin reagents

Cytoplasmic stains for standard HE staining

Eosin Y (yellowish eosin) solutions are commonly used as a counterstain to hematoxylin in the standard HE staining method. Eosin Y is an anionic dye that stains basic cell components (such as cytoplasm, collagen, muscle fibers and erythrocytes) bright red.

Eosin Y aqueous solutions

Eosin Y aqueous solutions for cytoplasmic counterstaining.

| | | |
|----------------------|--------------------|--------------------------------|
| Eosin Y 0.2% aqueous | 1000 mL 2500 mL | EOY-02-OT-1L EOY-02-OT-2.5L |
| Eosin Y 0.5% aqueous | 1000 mL 2500 mL | EOY-05-OT-1L EOY-05-OT-2.5L |
| Eosin Y 1% aqueous | 1000 mL 2500 mL | EOY-10-OT-1L EOY-10-OT-2.5L |
| Eosin Y 2% aqueous | 1000 mL 2500 mL | EOY-20-OT-1L EOY-20-OT-2.5L |

Eosin Y alcoholic solutions

Eosin Y alcoholic solutions for cytoplasmic counterstaining.

| | | |
|------------------------|--------------------|----------------------------------|
| Eosin Y 0.5% alcoholic | 1000 mL 2500 mL | EOYA-05-OT-1L EOYA-05-OT-2.5L |
| Eosin Y 1% alcoholic | 1000 mL 2500 mL | EOYA-10-OT-1L EOYA-10-OT-2.5L |
| Eosin Y 2% alcoholic | 1000 mL 2500 mL | EOYA-20-OT-1L EOYA-20-OT-2.5L |

Eosin Contrast

Modified alcoholic solution for cytoplasmic counterstaining. The reagent contains Eosin Y and Phloxine B for enhanced counterstaining effect.

| | |
|--------------------|----------------------------|
| 1000 mL 2500 mL | EOYK-OT-1L EOYK-OT-2.5L |
|--------------------|----------------------------|

Eosin Contrast PLUS

Modified alcoholic solution for intensive cytoplasmic counterstaining. Reagent contains Eosin Y, Phloxine B, and Biebrich Scarlet dyes for additional counterstaining effect.

| | |
|--------------------|------------------------------|
| 1000 mL 2500 mL | EOYKP-OT-1L EOYKP-OT-2.5L |
|--------------------|------------------------------|

Eosin Y instant

Mixture of powders for preparation of eosin reagent in aqueous or alcoholic medium.

| | | |
|----------------------------|-------------|-------------|
| Eosin Y Instant, aqueous | for 1000 mL | EOYI-AQ-1L |
| Eosin Y Instant, alcoholic | for 1000 mL | EOYI-ALC-1L |





Cytology and hematology staining reagents

Cytology and hematology staining kits contain reagents for fixation, staining and buffering cytology and hematology samples, as well as cytology smears of hematopoietic organs, providing precise distinction between cellular structures. Cytochemical staining kits contain reagents for cytochemical diagnosis of leukemia using bone marrow or whole blood smears, and also for staining and determining specific cell components in blood samples.

Papanicolaou (Pap) reagents

Nuclear and cytoplasmic reagents for Pap staining

Due to their simplicity of use and exceptional precision in diagnostics, staining reagents according to Papanicolaou have substantially contributed to decreasing occurrence of cervical cancer. By using a simple staining protocol, BioGnost's Pap reagents enable excellent differentiation between cellular structures.



Hematoxylin HP, Pap 1A

Modified Harris hematoxylin solution for nuclear staining according to Papanicolaou. Strong intensity reagent for progressive and regressive staining in exfoliative cytology.

| | |
|---------|---------------|
| 100 mL | HEMHP-OT-100 |
| 500 mL | HEMHP-OT-500 |
| 1000 mL | HEMHP-OT-1L |
| 2500 mL | HEMHP-OT-2.5L |

Hematoxylin HP, Pap 1B

Hematoxylin modified according to Harris for intensive nuclear staining according to Papanicolaou. Reagent for progressive and regressive staining in exfoliative cytology, with formulation stronger than Hematoxylin HP, Pap 1A.

| | |
|---------|-----------------|
| 100 mL | HEMHP1B-OT-100 |
| 500 mL | HEMHP1B-OT-500 |
| 1000 mL | HEMHP1B-OT-1L |
| 2500 mL | HEMHP1B-OT-2.5L |

OG-6 reagent, Pap 2A

Monochromatic reagent for counterstaining mature and keratinized squamous cell. Contains Orange G dye that stains the cytoplasm orange.

| | |
|---------|-------------|
| 100 mL | OG6-OT-100 |
| 500 mL | OG6-OT-500 |
| 1000 mL | OG6-OT-1L |
| 2500 mL | OG6-OT-2.5L |

Orange II reagent, Pap 2B

Monochromatic reagent for counterstaining mature and keratinized squamous cells. Contains Orange II dye that stains the cytoplasm intense red.

| | |
|---------|-------------|
| 100 mL | OR2-OT-100 |
| 500 mL | OR2-OT-500 |
| 1000 mL | OR2-OT-1L |
| 2500 mL | OR2-OT-2.5L |

EA 31 reagent, Pap 3A

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **gynecology** samples in cytology, contains the largest amount of green dye compared to other EA reagents. Stains the cytoplasm blue-green and pink, with emphasis on green and pink.

| | |
|---------|--------------|
| 100 mL | EA31-OT-100 |
| 500 mL | EA31-OT-500 |
| 1000 mL | EA31-OT-1L |
| 2500 mL | EA31-OT-2.5L |



EA 50 reagent, Pap 3B

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **gynecology** samples in cytology, stains the cytoplasm blue-green and pink, with emphasis on blue and pink.

| | |
|---------|--------------|
| 100 mL | EA50-OT-100 |
| 500 mL | EA50-OT-500 |
| 1000 mL | EA50-OT-1L |
| 2500 mL | EA50-OT-2.5L |



EA 65 reagent, Pap 3C

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **non-gynecology** samples in cytology, stains the cytoplasm red.

| | |
|---------|---------------|
| 100 mL | EA65C-OT-100 |
| 500 mL | EA65C-OT-500 |
| 1000 mL | EA65C-OT-1L |
| 2500 mL | EA65C-OT-2.5L |



EA 65 reagent, Pap 3D

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **non-gynecology** samples in cytology, stains the cytoplasm light blue-green and pink.

| | |
|---------|---------------|
| 100 mL | EA65D-OT-100 |
| 500 mL | EA65D-OT-500 |
| 1000 mL | EA65D-OT-1L |
| 2500 mL | EA65D-OT-2.5L |



Rapid Pap staining kit

Ready-to-use eight-reagent kit (in 20 containers that can be used as staining jars) for rapid progressive gynecology and non-gynecology cytological samples. Contains xylene substitute as clearing agent and xylene substitute-based medium for permanent covering of samples.

for 100 tests PAP-100T



OG-EA31 Pap reagent

One-step polychromatic reagent used as counterstain for gynecology samples in cytology. Enables rapid staining and differentiation between epithelial cells whose cytoplasm is stained green and pink. It is not necessary to use OG-6, Pap 2A reagent during staining because it stains cytoplasm of keratinized cell orange at the same time.

| | |
|---------|----------------|
| 100 mL | OGEA31-OT-100 |
| 500 mL | OGEA31-OT-500 |
| 1000 mL | OGEA31-OT-1L |
| 2500 mL | OGEA31-OT-2.5L |



OG-EA50 Pap reagent

One-step polychromatic reagent used as counterstain for gynecology samples in cytology. Enables rapid staining and differentiation between epithelial cells whose cytoplasm is stained blue and pink. It is not necessary to use OG-6, Pap 2A reagent during staining because it stains cytoplasm of keratinized cell orange at the same time.

| | |
|---------|----------------|
| 100 mL | OGEA50-OT-100 |
| 500 mL | OGEA50-OT-500 |
| 1000 mL | OGEA50-OT-1L |
| 2500 mL | OGEA50-OT-2.5L |



Shorr's solution

Enables clear differentiation of eosinophilic and cyanophilic cells of cervical epithelium. It is also used for a better visualization of sperm morphology.

| | |
|---------|------------|
| 100 mL | SH-OT-100 |
| 500 mL | SH-OT-500 |
| 1000 mL | SH-OT-1L |
| 2500 mL | SH-OT-2.5L |



Hematoxylin G1

Hematoxylin according to Gill, ideal for staining goblet cells. New generation reagent for delicate progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

| | |
|---------|---------------|
| 100 mL | HEMG1-OT-100 |
| 500 mL | HEMG1-OT-500 |
| 1000 mL | HEMG1-OT-1L |
| 2500 mL | HEMG1-OT-2.5L |

Hematoxylin G2


Hematoxylin according to Gill, contains double concentration of hematoxylin compared to Hematoxylin G1. Moderate intensity new generation reagent for progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

| | |
|---------|---------------|
| 100 mL | HEMG2-OT-100 |
| 500 mL | HEMG2-OT-500 |
| 1000 mL | HEMG2-OT-1L |
| 2500 mL | HEMG2-OT-2.5L |

Hematoxylin G3

Modified hematoxylin contains triple concentration of hematoxylin compared to Hematoxylin G1. Strong intensity new generation reagent for progressive staining in histopathology and cytology.

| | |
|---------|---------------|
| 100 mL | HEMG3-OT-100 |
| 500 mL | HEMG3-OT-500 |
| 1000 mL | HEMG3-OT-1L |
| 2500 mL | HEMG3-OT-2.5L |





Rapid and standard hematology reagents and kits

Bio-Diff kits

Rapid staining solutions for hematology, cytology and histology

Rapid diagnostic kits that provide precise staining results in a short period of time, such as results provided by May-Gruenwald Giemsa (MGG) staining method. Each component of Bio-Diff and Bio-Diff RTU kits is stabilized and prepared according to the highest standards.

Bio-Diff kit

Three-reagent kit that contains fixative agent, red and blue components for fast and effective staining. Each kit contains buffer tablets for consistent staining results.

| | |
|-----------|-----------|
| 3x100 mL | BD-K-100 |
| 3x500 mL | BD-K-500 |
| 3x1000 mL | BD-K-1L |
| 3x2500 mL | BD-K-2.5L |



Bio-Diff 1 reagent

Fixative agent, component of Bio-Diff kit.

| | |
|---------|-------------|
| 100 mL | BD1-OT-100 |
| 500 mL | BD1-OT-500 |
| 1000 mL | BD1-OT-1L |
| 2500 mL | BD1-OT-2.5L |



Bio-Diff 2 reagent

Buffered red component of Bio-Diff kit.

| | |
|---------|-------------|
| 100 mL | BD2-OT-100 |
| 500 mL | BD2-OT-500 |
| 1000 mL | BD2-OT-1L |
| 2500 mL | BD2-OT-2.5L |

Bio-Diff 3 reagent

Buffered blue component of Bio-Diff kit.

| | |
|---------|-------------|
| 100 mL | BD3-OT-100 |
| 500 mL | BD3-OT-500 |
| 1000 mL | BD3-OT-1L |
| 2500 mL | BD3-OT-2.5L |

STAINING DONE IN
14 SECONDS*



* Depending on the type of sample.

Bio-Diff RTU kit

Ready-to-Use three-reagent kit with reagents in containers that can be used as staining jars. Kit contains fixative agent, red and blue components for fast and effective staining and buffer tablet for consistent staining results.

3x100 mL BD-RTU-100



Bio-Diff 1 RTU reagent

Fixative component Bio-Diff RTU kit.

100 mL BD1-RTU



Bio-Diff 2 RTU reagent

Buffered red component of Bio-Diff RTU kit.

100 mL BD2-RTU

Bio-Diff 3 RTU reagent

Buffered blue component of Bio-Diff RTU kit.

100 mL BD3-RTU

STAINING DONE IN
14 SECONDS*



* depending on the type of sample

Buffer solution, pH 7.2

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

1000 mL PUF72-OT-1L
2500 mL PUF72-OT-2.5L
5000 mL PUF72-OT-5L

Buffer tablets pH 7.2

Phosphate buffer tablets for preparing buffer solutions used in hematology and histology. One buffer tablet is sufficient for preparation of 1000 mL of phosphate buffer solution.

50 pcs PT-72-50
100 pcs PT-72-100



Use buffer solution pH 6.8 for hematological and cytological diagnostics. Use buffer solution pH 7.2 for the detection of blood parasites.



May-Gruenwald and Giemsa reagents

Romanowsky staining solutions for hematology, cytology and histology

One of the most commonly used Pappenheim methods is May-Gruenwald Giemsa (MGG) for staining hematology and cytology materials, as well as cytology smears of hematopoietic organs. Giemsa solution is also used for staining histology sections.

May-Gruenwald solution

Polychromatic fixative solution containing Eosin Y, Methylene Blue and azure dyes. For staining in hematology, cytology and staining sections of hematopoietic organs in histopathology.

| | |
|-----------|------------|
| 100 mL | MG-OT-100 |
| 10x100 mL | MG-OT-110 |
| 500 mL | MG-OT-500 |
| 1000 mL | MG-OT-1L |
| 2500 mL | MG-OT-2.5L |



Giemsa solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For use in hematology, cytology, histology and for staining section samples of hematopoietic organs.

| | |
|-----------|------------|
| 100 mL | GM-OT-100 |
| 10x100 mL | GM-OT-110 |
| 500 mL | GM-OT-500 |
| 1000 mL | GM-OT-1L |
| 2500 mL | GM-OT-2.5L |



Buffer solution, pH 6.4

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

| | |
|---------|---------------|
| 1000 mL | PUF64-OT-1L |
| 2500 mL | PUF64-OT-2.5L |
| 5000 mL | PUF64-OT-5L |

Buffer solution, pH 6.8

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

| | |
|---------|---------------|
| 1000 mL | PUF68-OT-1L |
| 2500 mL | PUF68-OT-2.5L |
| 5000 mL | PUF68-OT-5L |

Buffer tablets pH 6.8

Phosphate buffer tablets for preparing solutions used in hematology and histology. One buffer tablet is sufficient for preparation of 1000 mL of phosphate buffer solution.

| | |
|---------|-----------|
| 50 pcs | PT-68-50 |
| 100 pcs | PT-68-100 |

Use buffer solution pH 6.8 for hematological and cytological diagnostics. Use buffer solution pH 7.2 for the detection of blood parasites.



Additional staining solutions for hematology and cytology



Wright's solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For staining in hematology, cytology and cytogenetics.

| | |
|---------|-----------|
| 100 mL | WR-OT-100 |
| 500 mL | WR-OT-500 |
| 1000 mL | WR-OT-1L |

Field kit

Ready to use two-reagent kit for rapid and efficient staining and detection of parasites in hematology samples. Kit for 100 tests contains reagents in containers that can be used as staining jars.

| | |
|---------------|----------|
| for 100 tests | FD-100T |
| 2x500 mL | FD-K-500 |

Wright-Giemsa solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. Modified Romanowsky stain for hematology staining.

| | |
|---------|--------------|
| 100 mL | WRGM-OT-100 |
| 500 mL | WRGM-OT-500 |
| 1000 mL | WRGM-OT-1L |
| 2500 mL | WRGM-OT-2.5L |



Hayem's solution

Solution for manual counting of erythrocytes.

| | |
|---------|-----------|
| 100 mL | HY-OT-100 |
| 500 mL | HY-OT-500 |
| 1000 mL | HY-OT-1L |

Tuerk's solution

Solution for manual counting of leukocytes.

| | |
|---------|-----------|
| 100 mL | TU-OT-100 |
| 500 mL | TU-OT-500 |
| 1000 mL | TU-OT-1L |

Brilliant Cresyl Blue, solution

Solution for staining reticulocytes in hematology.

| | |
|---------|------------|
| 100 mL | BCB-OT-100 |
| 500 mL | BCB-OT-500 |
| 1000 mL | BCB-OT-1L |

Leishman's solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For staining in hematology and clinical cytology.

| | |
|---------|-----------|
| 100 mL | LE-OT-100 |
| 500 mL | LE-OT-500 |
| 1000 mL | LE-OT-1L |

CytoGnost kit

Three-reagent kit for manual counting of human blood components.

| | |
|-------------|-----------|
| 3 x 100 mL | HCG-K-100 |
| 3 x 500 mL | HCG-K-500 |
| 3 x 1000 mL | HCG-K-1L |

ThromboGnost solution

Solution for manual counting of thrombocytes.

| | |
|---------|------------|
| 100 mL | THR-OT-100 |
| 500 mL | THR-OT-500 |
| 1000 mL | THR-OT-1L |



LEUKOGNOST

Cytochemical staining kits and ancillary reagents

Cytochemical staining kits

Enzymocytochemical blood smear and bone marrow staining of sections used to determine location and activity of cellular structures and enzyme systems in leukemia cell cytoplasm. Known as a routine method for diagnosis of hematological diseases and leukemia classification.

LeukoGnost SPE

Kit for detection of specific esterase activity (Leder stain, chloroacetate esterase) in leukocytes. Can be applied on formalin-fixed, paraffin-embedded tissue.

for at least 100 tests LKG-SPE



LeukoGnost ALP

Kit for detection of alkaline phosphatase activity in leukocytes.

for at least 100 tests LKG-ALP



LeukoGnost MPO

Kit for detection of myeloperoxidase activity in leukocytes.

for at least 100 tests LKG-MPO



LeukoGnost PAS

Kit for detection of the periodic acid-Schiff reaction in leukocytes.

for at least 100 tests LKG-PAS



LeukoGnost NSE

Kit for detection of non-specific esterase activity in leukocytes.

for at least 100 tests LKG-NSE



LeukoGnost SPENSE

Kit for simultaneous detection of specific and non-specific esterase activity in leukocytes.

for at least 50 tests LKG-SPENSE



LeukoGnost ACP

Kit for detection of alkaline phosphatase activity in leukocytes.

for at least 50 tests LKG-ACP



LeukoGnost PERLS

LeukoGnost PERLS (Prussian blue) kit for detection of free ferric (Fe³⁺) ions in blood and bone marrow. For application on tissue cells use HemoGnost PERLS kit.

for 100 tests LKG-PERLS

Sudan Black B kit

Four-reagent kit for staining neutrophil granules in hematological smears.

for 350-400 tests SBBH-K-100



Sudan Black B ECO

Phenol-free three-reagent kit for staining neutrophil granules in hematological smears.

for 100 tests SBBE-100T



LEUKOGNOST

Cytochemical staining kits and ancillary reagents

Ancillary reagents

Ancillary reagents greatly complement LeukoGnost kits - can be purchased separately or as a kit (LeukoGnost PLUS) containing a fixative, hematoxylin stain and a water-based mounting medium.

LeukoGnost PLUS

Additional set of reagents for use with LeukoGnost kits. Contains fixative, hematoxylin reagent and water-based mounting medium.

for at least 100 tests LKG-PLUS

LeukoGnost Fixative

Fixative for use in cytochemical diagnostics of leukemia

500 mL LKF-500

LeukoGnost Fixative ECO

Formalin-free fixative for use in cytochemical diagnostics of leukemia.

500 mL LFEC-500

LeukoGnost Fixative GL

Fixative for use in cytochemical diagnostics of leukemia, based on A formalin substitute.

500 mL LFGL-500

LeukoGnost HEM

Hematoxylin reagent for use with LeukoGnost kits.

| | |
|--------|------------|
| 100 mL | LKH-OT-100 |
| 250 mL | LKH-OT-250 |
| 500 mL | LKH-OT-500 |

BioMount Aqua

Aqueous covering medium for microscope sections and mounting medium for cover glasses.

| | |
|--------|---------|
| 30 mL | BMA-30 |
| 100 mL | BMA-100 |
| 250 mL | BMA-250 |
| 500 mL | BMA-500 |





Bacteriology and mycology staining reagents

Reagents for rapid and easy staining in microbiology allowing optimal differential analysis of bacteria. Phenol-free staining kits offer environment-friendly alternative without compromising quality of staining. Reagents for fungal analysis enable visualization of fungal structures (otherwise difficult to observe) and samples preservation.

BioGram kits and reagents

Kits for identification of bacteria according to Gram

Differential staining of bacteria according to Gram is widely known and used method in microbiology. BioGram 4 kit uses standard method to provide optimal staining and identification of Gram-positive and Gram-negative bacteria. BioGram ECO kit is a phenol-free, eco-friendly alternative to BioGram 4 kit with equal staining results.

BioGram 4 kit

Four-reagent kit for identification of bacteria according to Gram. Kit contains Gram Crystal Violet 1% solution, stabilized Gram Lugol solution, double amount of Gram Decolorizer solution 2 and Gram Safranin solution as counterstain.

| | |
|---------------|------------|
| for 100 tests | BGR4-100T |
| 5x100 mL | BGR4-K-100 |
| 5x250 mL | BGR4-K-250 |
| 5x500 mL | BGR4-K-500 |

Gram Crystal Violet 1% solution

Primary stain reagent for Gram-positive bacteria. Crystal Violet is also known as Gentian Violet and Methyl Violet 10B.

| | |
|---------|-------------|
| 100 mL | GC1-OT-100 |
| 250 mL | GC1-OT-250 |
| 500 mL | GC1-OT-500 |
| 1000 mL | GC1-OT-1L |
| 2500 mL | GC1-OT-2.5L |

Gram Lugol solution, stabilized

Solution contains polyvinylpyrrolidone iodine (PVP-I).

| | |
|---------|-------------|
| 100 mL | GLS-OT-100 |
| 250 mL | GLS-OT-250 |
| 500 mL | GLS-OT-500 |
| 1000 mL | GLS-OT-1L |
| 2500 mL | GLS-OT-2.5L |

Gram Decolorizer solution 2

Medium speed decolorizer used after stabilized iodine solution.

| | |
|---------|-------------|
| 100 mL | GD2-OT-100 |
| 250 mL | GD2-OT-250 |
| 500 mL | GD2-OT-500 |
| 1000 mL | GD2-OT-1L |
| 2500 mL | GD2-OT-2.5L |

Gram Safranin solution

Counterstain for Gram-negative bacteria, contains Safranin O dye.

| | |
|---------|-------------|
| 100 mL | GSF-OT-100 |
| 250 mL | GSF-OT-250 |
| 500 mL | GSF-OT-500 |
| 1000 mL | GSF-OT-1L |
| 2500 mL | GSF-OT-2.5L |

BioGram ECO kit PHENOL-FREE

Four-reagent phenol-free kit for identification of bacteria according to Gram. Kit contains Gram Crystal violet, phenol-free reagent, Gram Sodium hydrogencarbonate, solution, stabilized Gram Lugol solution, double amount of Gram Decolorizer solution 2 and Gram Safranin solution as counterstain.

| | |
|-------------------|------------|
| 2x50 mL+4x100 mL | BGRE-K-100 |
| 2x250 mL+4x500 mL | BGRE-K-500 |

Gram Crystal Violet, phenol-free reagent

Primary stain reagent for Gram-positive bacteria, doesn't contain phenol. Crystal Violet is also known as Gentian Violet and Methyl Violet 10B.

| | |
|---------|-------------|
| 100 mL | GCF-OT-100 |
| 250 mL | GCF-OT-250 |
| 500 mL | GCF-OT-500 |
| 1000 mL | GCF-OT-1L |
| 2500 mL | GCF-OT-2.5L |

Gram Sodium hydrogencarbonate, solution

Solution of sodium hydrogencarbonate for use with BioGram ECO kit.

| | |
|---------|--------------|
| 100 mL | GNHK-OT-100 |
| 250 mL | GNHK-OT-250 |
| 500 mL | GNHK-OT-500 |
| 1000 mL | GNHK-OT-1L |
| 2500 mL | GNHK-OT-2.5L |

Additional bacteriology staining solutions according to Gram

Gram Lugol solution

Non-stabilized Lugol solution.

| | |
|---------|-----------|
| 100 mL | GL-OT-100 |
| 250 mL | GL-OT-250 |
| 500 mL | GL-OT-500 |
| 1000 mL | GL-OT-1L |

Gram Decolorizer solution 1

Slow acting decolorizer solution used after non-stabilized iodine solution.

| | |
|---------|------------|
| 100 mL | GD1-OT-100 |
| 250 mL | GD1-OT-250 |
| 500 mL | GD1-OT-500 |
| 1000 mL | GD1-OT-1L |

Gram Decolorizer solution 3

Fast acting solution used with automated stainers.

| | |
|---------|------------|
| 100 mL | GD3-OT-100 |
| 250 mL | GD3-OT-250 |
| 500 mL | GD3-OT-500 |
| 1000 mL | GD3-OT-1L |

Gram Safranin 5x concentrated

Concentrated counterstain solution of Safranin O dye.

| | | |
|--|---------|--------------|
| For preparation of 500 mL of Safranin O counterstain solution | 100 mL | SO05X-OT-100 |
| For preparation of 2500 mL of Safranin O counterstain solution | 500 mL | SO05X-OT-500 |
| For preparation of 5000 mL of Safranin O counterstain solution | 1000 mL | SO05X-OT-1L |

Carbol Fuchsin Gram 10x concentrated

Concentrated Carbol Fuchsin solution for preparation of counterstain solution for modified staining. Also known as concentrated Gram Fuchsin solution.

| | | |
|--|---------|---------------|
| For preparation of 1000 mL of Carbol Fuchsin Gram counterstain solution | 100 mL | CFGR10-OT-100 |
| For preparation of 5000 mL of Carbol Fuchsin Gram counterstain solution | 500 mL | CFGR10-OT-500 |
| For preparation of 10000 mL of Carbol Fuchsin Gram counterstain solution | 1000 mL | CFGR10-OT-1L |

Additional iodine solutions

Lugol's solution

Classic formulation of aqueous solution of iodine and potassium iodide.

| | |
|--------|------------|
| 100 mL | LUG-OT-100 |
| 250 mL | LUG-OT-250 |
| 500 mL | LUG-OT-500 |

Lugol's C solution

Aqueous solution of iodine and potassium iodide, enhanced formulation with stronger staining (compared to classic Lugol solution).

| | |
|--------|-------------|
| 100 mL | LUGC-OT-100 |
| 250 mL | LUGC-OT-250 |
| 500 mL | LUGC-OT-500 |

Lugol's solution, acid

Aqueous iodine and potassium iodide solution with added acid.

| | |
|--------|-------------|
| 100 mL | LUGK-OT-100 |
| 250 mL | LUGK-OT-250 |
| 500 mL | LUGK-OT-500 |



TB-Stain kits and reagents

Kits for staining acid-fast bacteria according to Ziehl-Neelsen

Most commonly known method for detecting TB bacteria according to Ziehl-Neelsen. This method enables rapid and easy staining of acid-fast bacterial strains, such as *Mycobacteria*, *Nocardia*, *Rhodococcus*, *Dietzia* and *Gordonia*. TB-Stain ECO kit is a phenol-free, environment-friendly alternative to TB-Stain Hot kit.

TB-Stain Hot kit

Three-reagent kit for staining acid-fast bacteria. Contains TB Carbol Fuchsin reagent, double amount of TB Decolorizer and Methylene Blue Loeffler's reagent as counterstain.

| | |
|---------------|-----------|
| for 100 tests | TBH-100T |
| 4x100 mL | TBH-K-100 |
| 4x250 mL | TBH-K-250 |
| 4x500 mL | TBH-K-500 |

TB Carbol Fuchsin reagent

Synonym: Ziehl-Neelsen reagent. Primary stain reagent for acid-fast bacteria used in TB-Stain Hot, TB-Stain Cold and TB-Stain Quick kits.

| | |
|---------|-------------|
| 100 mL | TBC-OT-100 |
| 250 mL | TBC-OT-250 |
| 500 mL | TBC-OT-500 |
| 1000 mL | TBC-OT-1L |
| 2500 mL | TBC-OT-2.5L |

TB-Stain ECO kit

PHENOL-FREE

Three-reagent phenol-free kit for staining acid-fast bacteria. Contains TB-Fuchsin reagent, double amount of TB Decolorizer and Methylene Blue Loeffler's reagent as counterstain.

| | |
|----------|-----------|
| 4x100 mL | TBE-K-100 |
| 4x250 mL | TBE-K-250 |
| 4x500 mL | TBE-K-500 |

TB Decolorizer

For use in TB-Stain Hot and TB-Stain Cold kits.

| | |
|---------|-------------|
| 100 mL | TBD-OT-100 |
| 250 mL | TBD-OT-250 |
| 500 mL | TBD-OT-500 |
| 1000 mL | TBD-OT-1L |
| 2500 mL | TBD-OT-2.5L |

Methylene Blue Loeffler's reagent

Blue counterstain for TB-Stain Hot kit.

| | |
|---------|-------------|
| 100 mL | MBL-OT-100 |
| 250 mL | MBL-OT-250 |
| 500 mL | MBL-OT-500 |
| 1000 mL | MBL-OT-1L |
| 2500 mL | MBL-OT-2.5L |

TB Fuchsin reagent

Primary stain phenol-free reagent for acid-fast bacteria used in TB-Stain ECO kit.

| | |
|---------|--------------|
| 100 mL | TBFR-OT-100 |
| 250 mL | TBFR-OT-250 |
| 500 mL | TBFR-OT-500 |
| 1000 mL | TBFR-OT-1L |
| 2500 mL | TBFR-OT-2.5L |

Additional TB-Stain Hot kit staining solution

Methylene Blue 10x concentrated

Concentrated counterstain solution of Methylene Blue dye for use in microbiology.

| | | |
|--|--------|--------------|
| For preparation of 1000 mL of Methylene Blue counterstain solution | 100 mL | MB10X-OT-100 |
|--|--------|--------------|

| | | |
|--|--------|--------------|
| For preparation of 5000 mL of Methylene Blue counterstain solution | 500 mL | MB10X-OT-500 |
|--|--------|--------------|

| | | |
|---|---------|-------------|
| For preparation of 10000 mL of Methylene Blue counterstain solution | 1000 mL | MB10X-OT-1L |
|---|---------|-------------|

Kits for staining acid-fast bacteria according to Kinyoun and Kinyoun-Gabbett

TB-Stain Cold kit

Three-reagent kit for staining acid-fast bacteria according to Kinyoun. Contains TB Carbol Fuchsin reagent, double amount of TB Decolorizer and TB Malachite Green reagent as counterstain.

| | |
|---------------|-----------|
| for 100 tests | TBC-100T |
| 4x100 mL | TBC-K-100 |
| 4x250 mL | TBC-K-250 |
| 4x500 mL | TBC-K-500 |



TB Carbol Fuchsin reagent

Synonym: Ziehl-Neelsen reagent. Primary stain reagent for acid-fast bacteria used in TB-Stain Hot, TB-Stain Cold and TB-Stain Quick kits.

| | |
|---------|-------------|
| 100 mL | TBC-OT-100 |
| 250 mL | TBC-OT-250 |
| 500 mL | TBC-OT-500 |
| 1000 mL | TBC-OT-1L |
| 2500 mL | TBC-OT-2.5L |



TB Decolorizer

For use in TB-Stain Hot and TB-Stain Cold kits.

| | |
|---------|-------------|
| 100 mL | TBD-OT-100 |
| 250 mL | TBD-OT-250 |
| 500 mL | TBD-OT-500 |
| 1000 mL | TBD-OT-1L |
| 2500 mL | TBD-OT-2.5L |



TB Malachite Green reagent

Green counterstain for TB-Stain Cold kit.

| | |
|---------|-------------|
| 100 mL | TBM-OT-100 |
| 250 mL | TBM-OT-250 |
| 500 mL | TBM-OT-500 |
| 1000 mL | TBM-OT-1L |
| 2500 mL | TBM-OT-2.5L |

TB-Stain Quick kit

Three-reagent kit for rapid staining of acid-fast bacteria using Kinyoun-Gabbett method. Contains TB Carbol Fuchsin reagent and TB Armand reagent as counterstain.

| | |
|----------|-----------|
| 3x100 mL | TBQ-K-100 |
| 3x250 mL | TBQ-K-250 |
| 3x500 mL | TBQ-K-500 |



TB Armand reagent

Blue counterstain for use in TB-Stain Quick kit.

| | |
|---------|--------------|
| 100 mL | TBAR-OT-100 |
| 250 mL | TBAR-OT-250 |
| 500 mL | TBAR-OT-500 |
| 1000 mL | TBAR-OT-1L |
| 2500 mL | TBAR-OT-2.5L |



Additional TB-Stain Cold kit staining solution

Gabbett reagent

Blue counterstain for staining acid-fast bacteria according to modified Kinyoun method.

| | |
|---------|------------|
| 100 mL | GAB-OT-100 |
| 250 mL | GAB-OT-250 |
| 500 mL | GAB-OT-500 |
| 1000 mL | GAB-OT-1L |





Kits for detection of acid-fast bacteria using fluorescence microscopy according to modified Truant method

TB-Stain Fluorescent kit

Three-reagent kit for detection of acid-fast bacteria using fluorescence microscopy. Contains TB Auramine-Rhodamine reagent, double amount of TB Decolorizer Fluorescent and TB Permanganate reagent as counterstain.

| | |
|---------------|-----------|
| for 100 tests | TBF-100T |
| 4x100 mL | TBF-K-100 |
| 4x250 mL | TBF-K-250 |
| 4x500 mL | TBF-K-500 |



TB-Stain Auramine O kit

Three-reagent kit for staining acid-fast bacteria using fluorescence method. Contains TB Auramine O reagent, double amount of TB Decolorizer Fluorescent and counterstain of TB Permanganate reagent.

| | |
|----------|------------|
| 4x100 mL | TBAO-K-100 |
| 4x250 mL | TBAO-K-250 |
| 4x500 mL | TBAO-K-500 |



TB Auramine O reagent

Primary stain for use in TB Stain Auramine O kit.

| | |
|---------|--------------|
| 100 mL | TBAO-OT-100 |
| 250 mL | TBAO-OT-250 |
| 500 mL | TBAO-OT-500 |
| 1000 mL | TBAO-OT-1L |
| 2500 mL | TBAO-OT-2.5L |

TB Auramine-Rhodamine reagent

Primary stain reagent for use in TB-Stain Fluorescent kit.

| | |
|---------|-------------|
| 100 mL | TBA-OT-100 |
| 250 mL | TBA-OT-250 |
| 500 mL | TBA-OT-500 |
| 1000 mL | TBA-OT-1L |
| 2500 mL | TBA-OT-2.5L |



TB Decolorizer Fluorescent

For use in TB-Stain Fluorescent and TB-Stain Auramine O kits.

| | |
|---------|-------------|
| 100 mL | TBF-OT-100 |
| 250 mL | TBF-OT-250 |
| 500 mL | TBF-OT-500 |
| 1000 mL | TBF-OT-1L |
| 2500 mL | TBF-OT-2.5L |



TB Permanganate reagent

Counterstain for use in TB-Stain Fluorescent kit and TB-Stain Auramine O kits.

| | |
|---------|-------------|
| 100 mL | TBP-OT-100 |
| 250 mL | TBP-OT-250 |
| 500 mL | TBP-OT-500 |
| 1000 mL | TBP-OT-1L |
| 2500 mL | TBP-OT-2.5L |



Solutions for staining and analysis of fungi

In order to detect and achieve better visualization of fungi and fungal structures that are otherwise difficult to observe, fungi can be stained using Lactophenol solutions. Using Myco-Mount medium they can also be stained and preserved as a permanent preparation.

Lactophenol Blue solution

For use in mycology for staining and analysis of fungi.

| | |
|--------|------------|
| 100 mL | LPB-OT-100 |
| 250 mL | LPB-OT-250 |



Lactophenol solution

Colorless solution for use in mycology and analysis of fungi.

| | |
|--------|-----------|
| 100 mL | LP-OT-100 |
| 250 mL | LP-OT-250 |



For detection and fungal staining of histology sections, please see page 44 for more details on Grocott kit.

Myco-Mount Blue

Medium for simultaneous staining of fungi blue and preparation of permanent slides.

| | |
|--------|----------|
| 30 mL | MYMB-30 |
| 100 mL | MYMB-100 |



Myco-Mount Red

Medium for simultaneous staining of fungi red and preparation of permanent slides.

| | |
|--------|----------|
| 30 mL | MYMR-30 |
| 100 mL | MYMR-100 |



Potassium hydroxide, solution for fungi

Solution for use in mycology for analysis and visualization of fungi.

| | |
|--------|-------------|
| 30 mL | KHOG-OT-30 |
| 100 mL | KHOG-OT-100 |





Special staining kits for histology and cytology

Special kits for histology and cytology enable clear visualization of specific structures and cellular components. All of the reagents are easy to use and precisely formulated to maximize usability and to provide optimal results for the user.

Classification of special staining kits according to use:

MUSCLE AND/OR CONNECTIVE TISSUE

- A.F.O.G kit
- Azan Trichrome kit
- Elastica-Van Gieson kit
- Gomori Trichrome kit
- H.B.F.P. kit
- Hematoxylin P.T.A. kit
- Hematoxylin W kit
- Ladewig's Trichrome kit
- Mallory Trichrome kit
- Martius Scarlet Blue (MSB) kit
- Masson Trichrome kit
- Masson-Goldner Trichrome kit
- Movat kit
- Orcein kit
- Paraldehyde Fuchsin kit
- P.A.S.M. / Jones kit
- Picro-Sirius Red kit
- Safranin O kit
- Reticulin kit
- Reticulin Contrast kit
- Van Gieson Trichrome kit
- Verhoeff kit
- Weigert-Van Gieson kit

ENZYME ACTIVITY

- LeukoGnost SPE

FUNGI, BACTERIA AND PARASITES

- Alcian Yellow Toluidine Blue kit
- BioGram Histo kit
- Field kit
- Giemsa HP kit
- Grocott kit
- M.I.F. kit
- TB-Stain Histo kit
- Warthin Starry kit

LIPIDS

- Oil Red O kit
- Sudan Black B Lipid kit

NUCLEI AND NUCLEIC ACIDS

- Feulgen kit

PIGMENTS AND MINERAL DEPOSITS

- Fouchet-Van Gieson kit
- Grimelius kit
- HemoGnost Perls kit
- Masson Fontana kit
- Orcein kit
- Perls-Van Gieson kit
- Rhodanine kit
- Von Kossa kit

CYTOLOGY

- Bio-Diff RTU kit
- Eosin and nigrosin vital kit
- Sperm-Diff RTU kit
- UriGnost S kit
- UriGnost SM kit

CARBOHYDRATES


- Alcian Blue - P.A.S. kit
- Alcian Blue pH 1.0 kit
- Alcian Blue pH 2.5 kit
- Colloidal Iron kit
- Congo Red Highman kit
- Congo Red Puchtler kit
- Mucicarmine kit
- P.A.S. kit
- P.A.S. Diastase kit

CRYOSTAT

- Gomori Trichrome kit
- Oil Red O kit
- Orcein kit
- Sudan Black B Lipid kit

CENTRAL NERVOUS SYSTEM


- Bielschowsky kit
- Luxol Fast Blue kit



Elastica-Van Gieson kit

Four-reagent kit for staining elastic fibers and differentiation between elastic tissue, collagen and other types of connective tissue. The rapid method enables a satisfactory result with shorter section staining time.


for 100 tests WVGB-100T
4x100 mL WVGB-K-100



Eosin and nigrosine vital kit

Two-reagent kit containing separate dyes for rapid detection of sperm vitality and simple visualization of dead and living sperm cells. The nigrosin stain provides dark background for easier recognition of both viable and non-viable spermatozoa.


2x30 mL ENVK-K-30
2x100 mL ENVK-K-100



Feulgen kit

Five-reagent DNA staining kit according to Feulgen. For use in semiquantitative DNA determination in histological and cytological samples. Specimen is first treated with hydrochloric acid creating an aldehyde group of DNA that can be visualized by Schiff (BioSchiff) reagent. This reaction is specific for nuclear DNA.


for 100 tests FE-100T
6x100 mL FE-K-100



Field kit

Ready-to-use two-reagent kit for rapid and efficient staining and detection of parasites in hematology samples. Primarily used for staining thin and thick blood smears (dense drop) for purpose of diagnosing blood parasites. Reagents are stored in containers that can be used as staining jars.


for 100 tests FD-100T
2x500 mL FD-K-500



Fouchet-Van Gieson kit

Three-reagent kit for visualization of bilirubin and collagen according to Kutllick. Bilirubin is a yellow-brown pigment, but changes to green due to oxidation induced by Fouchet reaction. Green bilirubin can easily be detected on yellowish and pink colored background.

for 100 tests FVG-100T
3x100 mL FVG-K-100




Giemsa HP kit

Four-reagent kit for staining *Helicobacter pylori* in gastroscopic sections according to Lennart. Advantages of this method for detecting *H. pylori* are sensitive and reproducible results and easy performance.

for 100 tests GMHP-100T






Gomori Trichrome kit

Five-reagent kit for staining muscle, collagen fibre and nuclei, contains blue counterstain. Kit can be used to contrast skeletal, cardiac or smooth muscle.


for 100 tests GT-100T
5x100 mL GT-K-100
* Green counterstain also available



Grimelius kit

Five-reagent kit for staining argyrophilic granules. Grimelius kit can be used for detection of secretory intracytoplasmic granules specific for carcinoid tumors and for identification of neuroendocrine cells.

for 100 tests GRI-100T
5x100 mL GRI-K-100




Grocott kit, stabilized

Seven-reagent kit for visualization of fungi and histological argentaaffin structures in general (such as basal membranes). Green counterstain provides clear and visually rich contrast to target structures stained black.

for 100 tests GRC-100T
300-350 tests GRC-K-100


Stable at room temperature



H.B.F.P. kit

Three-reagent Hematoxylin-Basic Euchsin-Picric acid staining kit for detection of cardiac muscle changes after ischemia or myocardial infarction. H.B.F.P. kit is a non-enzymatic histochemical technique used for identification of early myocardial ischemia occurring on the cardiac muscle during the first six hours of symptoms.


for 100 tests HBFP-100T
3x100 mL HBFP-K-100



Hematoxylin P.T.A. kit

Four-reagent Hematoxylin-Phosphotungstic Acid kit for differentiation of smooth and striated muscle tissues as well as for detection of fibrin, collagen and elements of central nervous system according to Mallory.

for 100 tests HPTA-100T
4x100 mL HPTA-K-100




Hematoxylin W kit

Acid-resistant hematoxylin according to Weigert. Two-reagent kit that stains the nuclei blue-black, often a component of special staining kits for connective tissues.

for 100 tests HEMW-100T
2x100 mL HEMW-K-100
2x500 mL HEMW-K-500
2x1000 mL HEMW-K-1L






HemoGnost Perls kit

Three-reagent HemoGnost Perls (Prussian blue / Berlin blue) kit for the detection of reactive ferric (Fe³⁺) ions in tissue cells (like spleen or endometrial tissue). When staining bone marrow or other hematology tissue, use LeukoGnost PERLS kit (page 32).


for 100 tests HP-100T
3x250 mL HP-K-250



Ladewig Trichrome kit

Four-reagent kit for staining muscle (brown to red) and collagen fibers with blue counterstain. Staining procedure is easier and faster compared to Masson and Mallory Trichrome staining procedures.


for 100 tests LDW-100T
4x100 mL LDW-K-100



LeukoGnost SPE kit

Kit for detection of specific esterase activity (Leder stain, chloroacetate esterase) on formalin-fixed, paraffin-embedded (FFPE) samples of bone marrow tissue.

for at least 100 tests LKG-SPE



Luxol Fast Blue kit

Three-reagent kit for staining myelin and myelinated axons, Nissl bodies and phospholipids according to Kluver-Barrera. Kit is used for identifying the basic neuronal brain structure or spinal cord sections.


for 100 tests LFB-100T
3x100 mL LFB-K-100



M.I.F. kit

Two-reagent Merthiolate-Iodine-Formalin kit for fixing and staining fecal parasites (especially protozoae, cysts, helminth eggs and larvae). Stool sample is fixed with formalin and stained with two coloring agents; iodine and Eosin Y.


for 100 tests MIF-100T
for 200 tests MIF-200T



Mallory Trichrome kit

Three-reagent staining kit for connective tissue visualization and detection of collagen, cartilage, muscle, elastic fibers, mucous, pituitary cells, reticulum, bones, amyloid and erythrocytes.


for 100 tests MT-100T
3x100 mL MT-K-100



Martius Scarlet Blue (MSB) kit

Seven-reagent kit used for fibrin visualization, especially of older clusters. This method is a modification of Masson Trichrome and is ideal for studying connective tissue and vascular pathology.


for 100 tests MSB-100T
6x100 mL + 1x250 MSB-K-100
mL



Masson Fontana kit

Six-reagent melanine and argentaftin granule staining kit, based on the reduction of silver nitrate to elemental silver. Melanin is a brown-black pigment normally present in hair, skin, retina, iris and certain parts of CNS. Argentaftin granules are found in carcinoid tumors.


for 100 tests MF-100T
9x100 mL MF-K-100
9x500 mL MF-K-500



Masson-Goldner Trichrome kit

Seven-reagent kit for staining muscle and collagen fibers with green counterstain. Also used for visualizing gametes, nuclei, neurofibrils, glial cells, keratins, intercellular fibrils and for differentiation of smooth muscle fibers and collagens.


for 100 tests MGT-100T
7x100 mL MGT-K-100
7x500 mL MGT-K-500



Masson Trichrome kit

Seven-reagent kit for staining muscle and collagen fibers with blue counterstain. Also used for visualizing gametes, nuclei, neurofibrils, glial cells, keratins and intercellular fibrils. Kit may be useful for differentiation between collagen and smooth muscle fibers or detection of diseases like cirrhosis.


for 100 tests MST-100T
7x100 mL MST-K-100
7x500 mL MST-K-500



Movat kit

Eleven-reagent kit for staining collagen, elastic and muscle fibers, mucin and fibrin in tissue section. Movat kit is particularly useful when examining heart and vascular diseases.

for 100 tests MOV-100T
11x100 mL MOV-K-100




Mucicarmino kit

Mucicarmino kit is often used to identify primary tumor sites, distinguishing mucin-negative undifferentiated squamous cell lesions from mucin-positive adenocarcinomas. Can be also used for detection of diseases such as asthma, bronchitis, chronic obstructive pulmonary disease and cystic fibrosis.

for 100 tests MUC-100T
5x100 mL MUC-K-100






Oil Red O kit

Four-reagent kit for selective staining and detection of fat cells and neutral fats according to Johnson. Can be used with frozen sections and fresh smears to detect obesity-linked pathologies such as dyslipidemia and diabetes.


for 100 tests ORO-100T
4x250 mL ORO-K-250



Orcein kit

Five-reagent kit for visualization of hepatitis B surface antigen (HBsAg) seen as viral inclusion bodies in hepatocytes, for elastic fibers and copper associated protein in tissue sections. Can be used with frozen sections.


for 100 tests ORC-100T
5x250 mL ORC-K-250



Paraldehyde Fuchsin kit

Seven-reagent kit according to Gomori for detecting pathological changes in elastic fibers. It also stains mast cell granules, beta granules in pancreatic islets, neurosecretory material, mast cell granules and beta cells in the pituitary gland.


for 100 tests PAF-100T
9x100 mL PAF-K-100



P.A.S. kit

Five-reagent Periodic Acid-Schiff kit for staining aldehydes, mucopolysaccharides, mucoproteins and lymphocytes according to Hotchkiss-McManus. P.A.S. staining may also be used for the demonstration of fungal organisms in tissue sections.


for 100 tests PAS5-100T
5x100 mL PAS5-K-100
5x500 mL PAS5-K-500



P.A.S. Diastase kit

Five-reagent kit for detecting glycogen and mucopolysaccharide structures mostly in liver. The diastase digestion procedure, followed by P.A.S. staining, is useful in the evaluation of glycogen storage disease.

for 100 tests PDIA-100T




P.A.S.M. / Jones kit, stabilized

Seven-reagent Periodic Acid Silver Methenamine kit for staining kidney glomerular basement membranes. Kit includes red counterstain which provides visually rich contrast to target structures stained black.

for 100 tests PASM-100T
300-350 tests PASM-K-100


Stable at room temperature



Perls-Van Gieson kit

Three-reagent kit for the detection of reactive ferric (Fe^{3+}) ions in cells (Prussian blue/Berlin blue reaction) with Fuchsin Acid Van Gieson counterstain to differentiate collagen from connective tissue


for 100 tests PEG-100T
3x100 mL PEG-K-100



Picro-Sirius Red kit

Three-reagent kit for collagen staining and visualization in histology. Contains Sirius Red dye which enhances collagen birefringence so it can easily be identified by observing the section under polarized light.


for 100 tests PSR-100T



Reticulin kit

Seven-reagent kit for detecting argyrophilic reticulin fibers. It clearly differentiates between collagen and reticulin and nerve fibers and connective tissue. The main function of reticular fibers is to provide support and are normally found in liver, lymph nodes, spleen and kidneys.


for 100 tests RE-100T
7x50 mL RE-K-50
7x100 mL RE-K-100



Reticulin Contrast kit

Nine-reagent kit for detecting argyrophilic reticulin fibers according to Gordon and Sweets. The kit contains gold chloride solution that enhances visualization of reticulin fibers and it also contains Nuclear Fast Red (Kernechtrot) reagent that enables fine contrasting background.


for 100 tests RET-100T
9x50 mL RET-K-50
9x100 mL RET-K-100



Rhodanine kit

Four-reagent kit for detecting copper and copper-associated protein (CAP) in patients suffering from Wilson's disease. Abnormal copper accumulations are predominately found in liver tissue, but can also be found in the brain and cornea.


for 100 tests ROD-100T
5x100 mL ROD-K-100



Safranin O kit

Four-reagent kit for identification of cartilage, mucins and mastocyte granules used in histology and cytology.


for 100 tests SAFO-100T
4x100 mL SAFO-K-100



Sperm-Diff RTU kit

Ready-to-Use three-reagent kit with reagents stored in containers that can be used as staining jars. Kit contains fixative reagent, red and blue components for fast and effective staining.


3x100 mL SP-RTU-100



Sudan Black B Lipid kit

Four-reagent kit for specific lipid staining in cytochemistry. Sudan Black dye stains a few types of lipids, including neutral fats, phospholipids and sterols. Contains double amount of Sudan Black B, solution.


for 100 tests SBB-100T
6x100 mL SBB-K-100



TB-Stain Histo kit

Three-reagent kit for staining acid-fast bacteria (pathogenic mycobacteria) in histology sections, sputum, smears and culture smears according to Ziehl-Neelsen. Heating of the carbol-fuchsin solution is avoided in this protocol hence the release of hazardous phenolic vapors is omitted.

for 100 tests TBHK-100T



UriGnost S kit

Kit for sampling, staining, and microscopic analysis of urine sediment. Contains modified reagent according to Sternheimer (reagent manufactured according to the European Confederation of Laboratory Medicine (ECLM) and all the necessary equipment for sampling, concentrating, staining, and counting cells and kidney cylinders, as well as urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 µL pipette tips and urine plates).


for 500 tests USK-500
* for more information, see page 57



UriGnost SM kit

Kit for sampling, staining and microscopic analysis of urine sediments. Contains modified reagent according to Sternheimer-Malbin and equipment required for sampling, concentration, staining, counting of kidney cells and kidney cylinders and urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 µL pipette tips and urine plates).

for 500 tests USMK-500
* for more information, see page 57




Van Gieson Trichrome kit

Three-reagent kit for staining collagen fibers, muscle tissue, keratinized epithelium, cytoplasm, glial fibers and erythrocytes. Used for differentiation between collagen and smooth fibers in tumors and various other diseases.

for 100 tests VG-100T
3x100 mL VG-K-100
3x500 mL VG-K-500






Verhoeff kit

Six-reagent kit for detecting atrophy of elastic tissue in cases of emphysema, thinning and loss of elastic fibers in arteriosclerosis and other vascular diseases, or whether blood vessels have been invaded by a tumor.


for 100 tests VER-100T
6x100 mL VER-K-100



Von Kossa kit

Five-reagent kit for simple and reproducible detection of calcium deposits and calcium salt in tissue samples according to Von Kossa. Tissue calcification is associated with metabolic problems in various tissues (like bone marrow and mamma) and in tumors.


for 100 tests VK-100T
5x100 mL VK-K-100



Warthin Starry kit

Five-reagent kit for staining *Spirochaeta spp.*, *Helicobacter pylori*, *Microsporidia spp.* and *Legionella pneumophila*. The kit contains 12 jars with gelatin that enables both incubation and staining of sections, as well as other reagents that enable precipitation of silver on the bacterial surface. The bacteria are found in the mucus of the surface epithelium, in the apical gastric glands and in the gastric mucosa.

up to 48 tests WS-12T



Weigert-Van Gieson kit

Six-reagent kit for staining elastic fibers with prolonged incubation period. Used for differentiation between elastic tissue, collagen and other types of connective tissue.

for 100 tests WVG-100T
6x100 mL WVG-K-100



Additional staining reagents and solutions

Solutions and reagents (buffers, special stains, developer solutions and reagents for differentiation or counterstaining) that are either standalone reagents or components of special staining kits.

Staining reagents and solutions

4% Formaldehyde, alcoholic solution

For use with Reticulin and Reticulin Contrast staining kits.

| | |
|--------|------------|
| 100 mL | F4A-OT-100 |
|--------|------------|



Acetic buffer, Grimelius

For use with Grimelius kit

| | |
|--------|------------|
| 100 mL | APG-OT-100 |
|--------|------------|

Acid alcohols

For use with Azan Trichrome and Orcein staining kits.

| | | | |
|--------------------|---|--------|------------|
| Acid alcohol, Azan | ● | 100 mL | KAA-OT-100 |
| | | 250 mL | KAA-OT-250 |

| | | | |
|----------------------|---|--------|------------|
| Acid alcohol, Orcein | ● | 250 mL | KAO-OT-250 |
|----------------------|---|--------|------------|

Acid solution

For preparation of impregnation solution with Warthin Starry kit.

| | |
|--------|------------|
| 500 mL | KOT-OT-500 |
|--------|------------|

Acridine Orange solution

Acridine Orange aqueous solution is used as fluorescent dye for differential staining of DNA and RNA, acid mucins and detecting apoptosis; in cytogenetics for displaying structures containing abundant amounts of DNA. Also enables detection of microorganisms in blood smears.

| | |
|---------|------------|
| 100 mL | AOA-OT-100 |
| 500 mL | AOA-OT-500 |
| 1000 mL | AOA-OT-1L |

A.F.O.G. reagent

For use with A.F.O.G. (Acid Fuchsin Orange G) kit.

| | |
|--------|-------------|
| 100 mL | AFOG-OT-100 |
|--------|-------------|

Alcian Blue solution 1%

Alcian Blue 8GX dye 1% aqueous solution.

| | |
|--------|------------|
| 100 mL | AB1-OT-100 |
| 500 mL | AB1-OT-500 |

Alcian Blue solution pH 1.0

Reagent for staining heavily sulfated mucopolysaccharides. A component of Alcian Blue pH 1.0 kit.

| | |
|---------|-------------|
| 100 mL | AB10-OT-100 |
| 500 mL | AB10-OT-500 |
| 1000 mL | AB10-OT-1L |

Alcian Blue solution pH 2.5

Reagent for staining acid mucopolysaccharides of intestinal goblet cells. A component of Alcian Blue-P.A.S. and Alcian Blue pH 2.5 staining kits.

| | |
|---------|------------|
| 100 mL | AB2-OT-100 |
| 500 mL | AB2-OT-500 |
| 1000 mL | AB2-OT-1L |

Alkaline alcohol, solution

For use with Movat kit.

| | |
|--------|------------|
| 100 mL | ALA-OT-100 |
|--------|------------|





Additional staining reagents and solutions

Ammonium water

For use with Bielschowsky kit.

100 mL AV-OT-100

Ammonium iron sulfate, solution

For use with Reticulin and Reticulin Contrast staining kits.

100 mL ASF-OT-100

Aniline, alcoholic solution

For use with Azan Trichrome kit.

100 mL ANA-OT-100
250 mL ANA-OT-250

Aniline Blue reagent

Counterstain used with Masson Trichrome kit.

100 mL ABR-OT-100
500 mL ABR-OT-500

Aniline Blue, solution

For use with MSB kit.

100 mL ABO-OT-100

Azan reagent

For use with Azan Trichrome kit.

100 mL AZR-OT-100
250 mL AZR-OT-250

Azocarmine, solution

For use with Azan Trichrome kit.

100 mL AZ-OT-100
250 mL AZ-OT-250

Basic Activation buffer

For use with Oil Red O kit.

250 mL BAP-OT-250

Benedict's reagent

For detecting glucose in urine.

100 mL BNR-OT-100
250 mL BNR-OT-250
500 mL BNR-OT-500
1000 mL BNR-OT-1L

Biebrich Scarlet-Acid Fuchsin reagent

For use with special kits.

100 mL BSAF-OT-100
500 mL BSAF-OT-500

BioSchiff reagent

Schiff's reagent for detecting aldehydes and mucous substances. A component of several special staining kits.

100 mL BS-OT-100
500 mL BS-OT-500
1000 mL BS-OT-1L

BioSchiff Forte

Reagent for use with LeukoGnost PAS kit. Enhanced formulation of Schiff reagents for staining cytoplasm of neutrophils, monocytes, lymphocytes and basophils intensive magenta.

100 mL BSF-OT-100
500 mL BSF-OT-500

Borax, solution

For use with Grocott and P.A.S.M./Jones kits.

100 mL BO-OT-100
105 mL BO-OT-105
250 mL BO-OT-250

SCHIFF'S REAGENT
STABLE AT 22°C*



* In an unopened original packaging. Keep refrigerated after first use.

Chromotrope 2R / Aniline Blue reagent

Blue counterstain reagent, component of BioGnost's Gomori Trichrome kit.

100 mL CABR-OT-100

Chromotrope 2R / Fast Green F.C.F. reagent

Alternative green counterstain reagent, usually used with Gomori Trichrome kit.

100 mL CFG-OT-100

Colloidal Iron, stock solution

Reagent for binding ferric ions to acid mucopolysaccharides and proteoglycans.

100 mL KOL-OT-100

Congo Red Puchtler reagent

For use with Congo Red Puchtler kit.

100 mL CRPR-OT-100

Congo Red reagent

For use with Congo Red Highman kit.

100 mL CR-OT-100



Cresyl Violet, solution

For use with Luxol Fast Blue kit.

100 mL CV-OT-100

Developer stock solution

For use with Bielschowsky kit.

50 mL STR-OT-50

Differentiation reagents based on acetic acid

For use with special staining kits in histopathology.

| | | |
|---|--------|------------|
| Reagent for differentiation in Gomori Trichrome kit | 100 mL | RDG-OT-100 |
|---|--------|------------|

| | | |
|----------------------------|--------|-------------|
| Acetic acid, 0.5% solution | 100 mL | OK05-OT-100 |
|----------------------------|--------|-------------|

| | | |
|--------------------------|--------|------------|
| Acetic acid, 1% solution | 100 mL | OK1-OT-100 |
| | 500 mL | OK1-OT-500 |

| | | |
|--|--------|------------|
| Acetic acid, Colloidal Iron (A) solution ● | 250 mL | OKA-OT-250 |
|--|--------|------------|

| | | |
|--|--------|------------|
| Acetic acid, Colloidal Iron (B) solution | 100 mL | OKB-OT-100 |
|--|--------|------------|

Differentiation reagents based on hydrochloric acid

For use with special staining kits in histopathology.

| | | |
|-------------------------|--------|--------------|
| HCL reagent, A.F.O.G. ● | 100 mL | HCLAF-OT-100 |
|-------------------------|--------|--------------|

| | | |
|--------------------------|--------|--------------|
| HCL reagent, Alcian Blue | 100 mL | HCLAB-OT-100 |
|--------------------------|--------|--------------|

| | | |
|-------------------------------|--------|-------------|
| HCL reagent, Colloidal Iron ● | 100 mL | HCLK-OT-100 |
|-------------------------------|--------|-------------|

| | | |
|------------------------|--------|-------------|
| HCL reagent, Feulgen ● | 100 mL | HCLF-OT-100 |
|------------------------|--------|-------------|

| | | |
|------------------------------|--------|-------------|
| HCL reagent, HemoGnost Perls | 100 mL | HCLH-OT-100 |
|------------------------------|--------|-------------|

| | | |
|------------------------------|--------|-------------|
| HCL reagent, HemoGnost Perls | 250 mL | HCLH-OT-250 |
|------------------------------|--------|-------------|

| | | |
|---------------------|--------|-------------|
| HCL reagent, P.A.S. | 100 mL | HCLP-OT-100 |
|---------------------|--------|-------------|

| | | |
|---------------------|--------|-------------|
| HCL reagent, P.A.S. | 500 mL | HCLP-OT-500 |
|---------------------|--------|-------------|

| | | |
|-----------------------------------|--------|-------------|
| HCL reagent, Weigert-Van Gieson ● | 100 mL | HCLW-OT-100 |
|-----------------------------------|--------|-------------|

Ehrlich aldehyde reagent

Reagent for detecting urobilinogen in urine.

100 mL EAR-OT-100

Eosin-nigrosin vital

Fast detection (one-step detection) of sperm vitality and visualization of dead and living sperm cells with one reagent. Simple, easy and fast method for semen analysis.

30 mL ENV-30

100 mL ENV-100

Eosin Y, isotonic solution

Aqueous solution of Eosin Y used with Eosin and nigrosin vital kit.

100 mL EIO-OT-100

ErioGnost reagent

Acid fast reagent for nuclear staining, comparable to hematoxylin staining. Ecologically acceptable synthetic replacement for hematoxylin reagents.

100 mL ERS-OT-100

500 mL ERS-OT-500

1000 mL ERS-OT-1L

Fast Green F.C.F. reagent

Counterstain used with Masson-Goldner Trichrome and Paraldehyde Fuchsin staining kits.

100 mL FGR-OT-100

500 mL FGR-OT-500

Fast Green F.C.F. contrast reagent

Counterstain used in Grocott and Feulgen staining kits.

100 mL FGKR-OT-100

Field, solution A

Blue component of Field kit.

100 mL FDA-RTU

500 mL FDA-OT-500

Field, solution B

Red component of Field kit.

100 mL FDB-RTU

500 mL FDB-OT-500

Ferri reagent, Verhoeff B

For use with Verhoeff kit.

100 mL FRV-OT-100

Ferri reagent, Weigert B

Iron(III) nitrate solution in diluted hydrochloric acid, a component of Hematoxylin W kit.

100 mL FR-OT-100

500 mL FR-OT-500

1000 mL FR-OT-1L

Fuchsin Acid reagent

For use with Mallory Trichrome kit.

100 mL FA-OT-100



Additional staining reagents and solutions

Fuchsin Acid Van Gieson reagent

For use in special staining kits.
Synonym: Picrofuchsin reagent.

100 mL FAG-OT-100

Fuchsin Basic reagent

For use in bacteriology.

100 mL FBR-OT-100
500 mL FBR-OT-500
1000 mL FBR-OT-1L

Fuchsin Basic, solution

For use with H.B.F.P. kit.

100 mL FBO-OT-100

Gelatin, gel in transport jar

For preparing developing solution with Warthin Starry staining kit.

12x8 mL GEL-128



Glutaraldehyde, 4% buffered solution

Solution used with Sudan Black B kit.

100 mL G4-OT-100



Gold chloride solutions

Aqueous solutions of gold chloride (aurichloric acid, HAuCl_4), component of many special staining kits.

| | | |
|------------------------------|--------|-------------|
| Gold chloride, 0.2% solution | 100 mL | ZK02-OT-100 |
| Gold chloride, 0.6% solution | 100 mL | ZK06-OT-100 |
| Gold chloride, 1% solution | 50 mL | ZK1-OT-50 |
| Gold chloride, 2% solution | 50 mL | ZK2-OT-50 |

Hematoxylin, Verhoeff A

For use with Verhoeff and Movat kits.

100 mL HEMV-OT-100



Hematoxylin, Weigert A

Hematoxylin solution in ethanol, component of Hematoxylin W kit.
Part of several special staining kits.

100 mL HEMA-OT-100
500 mL HEMA-OT-500
1000 mL HEMA-OT-1L



Hydroquinone, solution

For preparation of developing solution in Warthin Starry kit.

50 mL HQ-OT-50



Hydroquinone-sodium sulphite

For use with Grimelius kit

15 x 3,5 g HNS-15-35

Iodine solution, Verhoeff C

For use with Verhoeff and Movat kits.

100 mL JODV-OT-100

Iron chloride, solution

For use with Fouchet-Van Gieson kit.

100 mL FK-OT-100

Kovac indole reagent

Reagent for identification of indole-positive and indole-negative microorganisms.

100 mL KOV-OT-100



Ladewig's reagent

For use with Ladewig Trichrome kit, contains Aniline Blue, Acid Fuchsin and Orange G dyes.

100 mL LDWR-OT-100

Light Green S.F. reagent

Counterstain used with trichrome special staining kits.

100 mL LGS-OT-100

Lithium carbonate, Luxol

Lithium carbonate solution used with Luxol Fast Blue kit.

100 mL LKL-OT-100

Lithium carbonate, solution

Lithium carbonate solution used with Von-Kossa staining kit.

100 mL LK-OT-100

Lugol's solution, M.I.F.

For use with M.I.F. kit.

100 mL LUGM-OT-100

Luxol Fast Blue, solution

For use with Luxol Fast Blue kit.

100 mL LFB-OT-100



M.F. reagent

For use with M.I.F. kit.

100 mL MF-OT-100

Martius Yellow, solution

For use with MSB kit.

100 mL MAY-OT-100

Mayer's mucicarmine

For use with Mucicarmine kit.

100 mL MMU-OT-100



Metanil Yellow, solution

For use with Mucicarmine kit.

100 mL MY-OT-100

Methenamine, solution

For use with Grocott and P.A.S.M./Jones kits, stabilized.

| | |
|--------|------------|
| 100 mL | MET-OT-100 |
| 250 mL | MET-OT-250 |

Methyl Blue Eosin, solution

Modified Mann's solution for use in histology.

| | |
|---------|------------|
| 100 mL | MBE-OT-100 |
| 500 mL | MBE-OT-500 |
| 1000 mL | MBE-OT-1L |

Neutral Red reagent

Neutral Red aqueous counterstain solution for use with special staining kits.

| | |
|--------|-----------|
| 100 mL | NR-OT-100 |
|--------|-----------|

Nigrosine, isotonic solution

Aqueous solution of nigrosine used with Eosin and nigrosine vital kit.

| | |
|--------|------------|
| 100 mL | NIO-OT-100 |
|--------|------------|

Nuclear Fast Red (Kernechtrot) reagent

Nuclear Fast Red dye solution for staining nuclei. Counterstain in many special staining kits.

| | |
|--------|-----------|
| 100 mL | KR-OT-100 |
| 250 mL | KR-OT-250 |
| 500 mL | KR-OT-500 |

Oil Red O reagent

Lipid substances staining solution.

| | |
|--------|------------|
| 250 mL | ORO-OT-250 |
|--------|------------|

Orange G, 1% solution

For use with Movat kit.

| | |
|--------|-------------|
| 100 mL | ORG1-OT-100 |
|--------|-------------|

Orange G / Aniline Blue reagent

For use with Mallory Trichrome kit.

| | |
|--------|------------|
| 100 mL | OGA-OT-100 |
|--------|------------|

Orcein reagent

For use with Orcein kit.

| | |
|--------|-------------|
| 250 mL | ORCR-OT-250 |
|--------|-------------|

Oxalic acid solutions

For use with several special staining kits.

| | | |
|--------------------------|--------|-------------|
| Oxalic acid, 1% solution | 100 mL | OKS1-OT-100 |
| Oxalic acid, 2% solution | 250 mL | OKS2-OT-250 |

Paraldehyde Fuchsin reagent

Reagent for staining elastins and mucins.

| | |
|--------|------------|
| 100 mL | PAF-OT-100 |
|--------|------------|

Periodic acid, solution

Synonym: Orthoperiodic acid. Oxidizing medium widely used in histology. Key component of many special staining kits, such as P.A.S. and Alcian Blue-P.A.S., P.A.S.M./Jones, stabilized, Grocott kit, stabilized and Weigert-Van Gieson kit.

| | | |
|------------------------------|--------|-------------|
| Periodic acid, 0.5% solution | 100 mL | PK05-OT-100 |
| Periodic acid, 0.5% solution | 250 mL | PK05-OT-250 |
| Periodic acid, 0.5% solution | 500 mL | PK05-OT-500 |

| | | |
|------------------------------|--------|-------------|
| Periodic acid, 0.8% solution | 100 mL | PK08-OT-100 |
| Periodic acid, 0.8% solution | 250 mL | PK08-OT-250 |
| Periodic acid, 0.8% solution | 500 mL | PK08-OT-500 |

| | | |
|----------------------------|--------|------------|
| Periodic acid, 1% solution | 100 mL | PK1-OT-100 |
| Periodic acid, 1% solution | 250 mL | PK1-OT-250 |
| Periodic acid, 1% solution | 500 mL | PK1-OT-500 |

Picro-Sirius Red reagent

For use with Picro-Sirius Red kit for collagen staining and differentiation.

| | |
|--------|------------|
| 100 mL | PSR-OT-100 |
|--------|------------|

Phosphomolybdic acid, 1% solution

For use with A.F.O.G. (Acid Fuchsin Orange G) and Mallory Trichrome staining kits.

| | |
|--------|-------------|
| 100 mL | FMK1-OT-100 |
|--------|-------------|

Phosphomolybdic acid, 5% solution

For use with Azan Trichrome kit.

| | |
|--------|-------------|
| 100 mL | FMK5-OT-100 |
| 250 mL | FMK5-OT-250 |

Phosphotungstic acid, 1% solution

For use with Martius Scarlet Blue (MSB) kit.

| | |
|--------|-------------|
| 100 mL | FVK1-OT-100 |
| 500 mL | FVK1-OT-500 |

Phosphotungstic acid, 5% solution

For use with Movat kit.

| | |
|--------|-------------|
| 100 mL | FVK5-OT-100 |
|--------|-------------|

Picric acid in acetone, solution

For use with H.B.F.P. and BioGram Histo kits.

| | |
|--------|------------|
| 100 mL | PKA-OT-100 |
|--------|------------|

Ponceau S, solution

For use with MSB kit.

| | |
|--------|-------------|
| 100 mL | PONS-OT-100 |
|--------|-------------|

Potassium hexacyanoferrate, Colloidal Iron solution

Potassium hexacyanoferrate (II) solution for use with Colloidal Iron kit.

| | |
|--------|-------------|
| 100 mL | KHCK-OT-100 |
|--------|-------------|



Additional staining reagents and solutions

Potassium hexacyanoferrate, solution

Potassium hexacyanoferrate (II) solution, optimally concentrated for use with HemoGnost Perls kit.

| | |
|--------|------------|
| 100 mL | KHC-OT-100 |
| 250 mL | KHC-OT-250 |

Potassium hydroxide, solution

Optimally concentrated aqueous solution of potassium hydroxide used with Congo Red Highman kit.

| | |
|--------|------------|
| 100 mL | KHO-OT-100 |
|--------|------------|

Potassium permanganate, solutions

For use with Hematoxylin P.T.A., Masson Fontana, Orcein, Paraldehyde Fuchsin and Reticulin staining kits.

| | | |
|---------------------------------------|--------|-------------|
| Potassium permanganate, 0.5% solution | 100 mL | KP05-OT-100 |
|---------------------------------------|--------|-------------|

| | | |
|-------------------------------------|--------|------------|
| Potassium permanganate, 1% solution | 250 mL | KP1-OT-250 |
|-------------------------------------|--------|------------|

Propylene glycol

For use with Sudan Black B Lipid kit.

| | |
|--------|--------|
| 100 mL | PG-100 |
|--------|--------|

Propylene glycol, 85%

For use with Sudan Black B Lipid kit.

| | |
|--------|----------|
| 100 mL | PG85-100 |
|--------|----------|

P.T.A. - P.M.A. reagent

Reagent containing optimal concentration of phosphotungstic and phosphomolybdic acid. Used with Masson Trichrome and Masson-Goldner Trichrome staining kits.

| | |
|--------|------------|
| 100 mL | PPR-OT-100 |
| 500 mL | PPR-OT-500 |

Reagent for differentiation in Verhoeff kit

For use with Verhoeff kit.

| | |
|--------|------------|
| 100 mL | RDV-OT-100 |
|--------|------------|

Reducing buffer, solution

For use with Von Kossa kit.

| | |
|--------|-----------|
| 100 mL | RP-OT-100 |
|--------|-----------|

Resorcine Fuchsin reagent

For use with Weigert-Van Gieson kit.

| | |
|--------|------------|
| 100 mL | RFR-OT-100 |
|--------|------------|

Rhodanine reagent

For use with Rhodanine kit.

| | |
|--------|-----------|
| 100 mL | RR-OT-100 |
|--------|-----------|

Safranin O reagent

Counterstaining reagent for use in histology and cytology. Used for visualizing and staining cartilage, mucins and mastocytes in formalin-fixed and paraffin-embedded tissue samples.

| | |
|--------|-------------|
| 100 mL | SAFO-OT-100 |
|--------|-------------|

Silver nitrate solutions

For use in special staining kits.

| | | |
|---|--------|-----------|
| Silver ammonia solution (component of Reticulin and Reticulin Contrast staining kits) ● | 100 mL | SA-OT-100 |
|---|--------|-----------|

| | | |
|--|--------|------------|
| Silver ammonia reagent (component of Von Kossa and Masson Fontana staining kits) ● | 100 mL | SAR-OT-100 |
|--|--------|------------|

| | | |
|--|--------|-------------|
| Silver ammonia reagent stock solution (component of Bielschowsky staining kit) ● | 100 mL | SSAR-OT-100 |
|--|--------|-------------|

| | | |
|---|--------|-------------|
| Silver nitrate, 20% solution (component of Bielschowsky staining kit) ● | 100 mL | SN20-OT-100 |
|---|--------|-------------|

| | | |
|---|--------|------------|
| Silver nitrate solution, stabilized (component of Grocott and P.A.S.M. / Jones, stabilized staining kits) | 100 mL | SNS-OT-100 |
|---|--------|------------|

| | | |
|--|--------|-------------|
| Silver nitrate, WS solution (component of Warthin Starry staining kit) ● | 100 mL | SNWS-OT-100 |
|--|--------|-------------|

Sodium acetate, solution

For use with Rhodanine kit.

| | |
|--------|-----------|
| 100 mL | NA-OT-100 |
|--------|-----------|

Sodium hydroxide, solution

For use with Congo Red Puchtler kit.

| | |
|-------|----------|
| 30 mL | NH-OT-30 |
|-------|----------|

Sodium metabisulfite solutions

Component of Alcian Blue-P.A.S., P.A.S. and Feulgen staining kits.

| | | |
|----------------------|--------|-----------|
| Sodium metabisulfite | 100 mL | NM-OT-100 |
| | 500 mL | NM-OT-500 |

| | | |
|--|--------|------------|
| Sodium metabisulfite, solution for Feulgen | 100 mL | NMF-OT-100 |
|--|--------|------------|

Sodium tetraborate, solution

A component of Alcian Blue pH 2.5 and Alcian Blue-P.A.S. staining kits.

| | |
|--------|------------|
| 100 mL | NTB-OT-100 |
| 500 mL | NTB-OT-500 |

Sodium thiosulfate solution

Solutions are component of Grocott stabilized, Masson Fontana, P.A.S.M. / Jones, stabilized, Reticulin, Reticulin Contrast and Von Kossa staining kits.

| | | |
|---------------------------------|--------|------------|
| Sodium thiosulfate, 2% solution | 100 mL | NT2-OT-100 |
|---------------------------------|--------|------------|

| | | |
|---------------------------------|--------|------------|
| Sodium thiosulfate, 5% solution | 100 mL | NT5-OT-100 |
|---------------------------------|--------|------------|

Sudan Black B, solution

Solution for staining lipids, used with Sudan Black B lipid kit.

| | |
|--------|------------|
| 100 mL | SBO-OT-100 |
|--------|------------|

Sudan Black B reagent

Staining reagent for use in hematology, used with Sudan Black B kit.

100 mL SBB-OT-100



Sulfosalicylic acid, 20% solution

Reagent for detecting proteins in urine.

100 mL SSK-OT-100

Sulfuric acid solutions

Used with Orcein, Paraldehyde Fuchsin and Reticulin staining kits.

| | | |
|------------------------------|--------|-------------|
| Sulfuric acid, 0.3% solution | 250 mL | SK03-OT-250 |
|------------------------------|--------|-------------|

| | | |
|------------------------------|--------|-------------|
| Sulfuric acid, 0.5% solution | 100 mL | SK05-OT-100 |
|------------------------------|--------|-------------|

| | | |
|----------------------------|--------|------------|
| Sulfuric acid, 3% solution | 100 mL | SK3-OT-100 |
|----------------------------|--------|------------|

Trichloroacetic acid, solution

For use with Fouchet-Van Gieson kit.

100 mL TKO-OT-100



Toluidine Blue, solution

Solution for nuclear staining and metachromatic staining of mastocytes in connective tissue.

100 mL TB-OT-100

250 mL TB-OT-250

UriGnost S reagent

Modification according to Sternheimer for staining and microscopical analysis of urine sediment. A component of UriGnost S kit.

| | | |
|----------------------------------|-------|---------|
| UriGnost S reagent for 500 tests | 50 mL | UGS-500 |
|----------------------------------|-------|---------|

| | | |
|-----------------------------------|--------|----------|
| UriGnost S reagent for 1000 tests | 100 mL | UGS-1000 |
|-----------------------------------|--------|----------|

* for more information, see page 49

UriGnost SM reagent

Modification according to Sternheimer-Malbin for staining and microscopical analysis of urine sediment. A component of UriGnost SM kit.

| | | |
|-----------------------------------|-------|----------|
| UriGnost SM reagent for 500 tests | 50 mL | UGSM-500 |
|-----------------------------------|-------|----------|

| | | |
|------------------------------------|--------|-----------|
| UriGnost SM reagent for 1000 tests | 100 mL | UGSM-1000 |
|------------------------------------|--------|-----------|

* for more information, see page 49



Azure II Eosinate

Azure II eosinate. For preparation of Romanowsky stains, made by mixing equal ratios of Azure B, Methylene Blue and Eosin Y stains.

| | |
|------|----------|
| 25 g | A2E-P-25 |
|------|----------|

Biebrich Scarlet, C.I. 26905

Croceine Scarlet, Ponceau B, Ponceau BS, BSC certified stain. For Masson trichrome staining and a component of Shorr's solution.

| | |
|------|---------|
| 25 g | BS-P-25 |
|------|---------|

Bismarck Brown Y, C.I. 21000

Bismarck Brown Y, Vesuvine BA, BSC certified. For use in histopathology and cytology.

| | |
|------|---------|
| 25 g | BB-P-25 |
|------|---------|

Brilliant Cresyl Blue, C.I. 51010

For use in hematology (for reticulocyte staining) and histology (for acid mucopolysaccharides).

| | |
|------|----------|
| 10 g | BCB-P-10 |
| 25 g | BCB-P-25 |

Brilliant Green Certified, C.I. 42040

Malachite Green G, Basic Green 1, Emerald Green. For bacteriostatic use.

| | |
|------|---------|
| 25 g | BG-P-25 |
|------|---------|

Carmine, C.I. 75470

Natural Red 4, Cochineal, Nacaral, carminic acid and aluminum compound. For staining mucins and glycogens.

| | |
|------|----------|
| 5 g | CAR-P-5 |
| 10 g | CAR-P-10 |
| 25 g | CAR-P-25 |

Carminic acid, C.I. 75470

Natural Red 4. For staining mucines and glycogen.

| | |
|------|-----------|
| 5 g | CARA-P-5 |
| 10 g | CARA-P-10 |
| 25 g | CARA-P-25 |

Celestine Blue, C.I. 51050

Celestine blue B, Mordant blue 14. For staining nuclei dark blue, used as hematoxylin substitute.

| | |
|------|---------|
| 10 g | CB-P-10 |
| 25 g | CB-P-25 |

Chromotrope 2R, C.I. 16570

Acid Red 29, Mordant Blue 80. For counterstaining cytoplasm in histopathology used as eosin substitute in HE staining and in trichrome stains.

| | |
|------|----------|
| 25 g | C2R-P-25 |
|------|----------|

Congo Red, C.I. 22120

Direct Red 28, Kongorot, BSC certified stain. For staining of amyloids according to Highman and Puchtler and for elastic fiber staining.

| | |
|------|---------|
| 25 g | CR-P-25 |
|------|---------|

Cresyl Violet Acetate Certified

Cresyl Violet, Oxazine 9, Cresyl Echt Violet. For use in microbiology and histology.

| | |
|------|----------|
| 25 g | CVA-P-25 |
|------|----------|

Crystal Violet, C.I. 42555

Gentian Violet, Methyl Violet 10B, BSC certified stain. For bacterial staining according to Gram.

| | |
|------|----------|
| 25 g | CRV-P-25 |
| 50 g | CRV-P-50 |

Eosin B, C.I. 45400

Eosin B, Eosin Bluish, Imperial Red, Acid Red 91, BSC certified. For staining according to Romanowsky. Used for staining red blood cells, connective tissues and as counterstain for the Gram stain.

| | |
|------|---------|
| 25 g | EB-P-25 |
|------|---------|

Eosin Y, C.I. 45380

Eosin Yellowish, Acid Red 87, Eosin WS, BSC certified stain. Counterstain for hematoxylin and for preparing EA Pap stains and polychromatic hematological stains.

| | |
|-------|-----------|
| 25 g | EOY-P-25 |
| 100 g | EOY-P-100 |
| 500 g | EOY-P-500 |

Eriochrome Cyanine R, C.I. 43820

Mordant Blue 3, Chromoxane Cyanine R. For use in cytology and histology.

| | |
|------|----------|
| 25 g | ECR-P-25 |
|------|----------|

Fast Green F.C.F., C.I. 42053

Food Green 3, Solid Green F.C.F, 1724 Green, BSC certified stain. For trichrome staining according to Gomori and Masson.

| | |
|------|---------|
| 25 g | FG-P-25 |
|------|---------|

Fuchsin Acid, C.I. 42685

Acid fuchsine, Acid Violet 19, BSC certified stain. For staining connective tissue according to Van Gieson and Mallory.

| | |
|------|---------|
| 25 g | FA-P-25 |
|------|---------|

Fuchsin Basic, C.I. 42510

Basic fuchsine, Basic Violet 14, BSC certified stain. For staining mucins, Negri bodies and acid-fast bacteria.

| | |
|-------|----------|
| 25 g | FB-P-25 |
| 100 g | FB-P-100 |

Fuchsin Basic - Schiff, C.I. 42500

Pararosaniline HCl, Basic Red 9, BSC certified stain. Used for P.A.S. (Periodic Acid Schiff) staining, DNA staining according to Feulgen and acid-fast bacteria staining.

| | |
|------|----------|
| 25 g | FBS-P-25 |
| 50 g | FBS-P-50 |

Gentian Violet, C.I. 42555

Crystal Violet, Methyl Violet 10B, BSC certified stain. For bacterial staining according to Gram.

| | |
|-------|----------|
| 25 g | GV-P-25 |
| 100 g | GV-P-100 |

Giemsa

BSC certified stain. Romanowsky dye for staining according to Pappenheim. Polychromatic dye used for identification of chromosomal damage and as a differentiating stain.

| | |
|------|--------|
| 25 g | G-P-25 |
| 50 g | G-P-50 |



Hematoxylin, C.I. 75290

Natural Black 1, BSC certified stain. For preparing nuclear staining reagents.

| | |
|-------|---------|
| 25 g | H-P-25 |
| 100 g | H-P-100 |

Janus Green B, C.I. 11050

Diazine Green, BSC certified stain. For displaying mitochondria in living cells.

| | |
|------|---------|
| 25 g | JG-P-25 |
|------|---------|

Jenner Stain

Jenner, BSC certified stain. Romanowsky polychromatic stain routinely used in hematology for staining blood smears.

| | |
|------|---------|
| 25 g | JS-P-25 |
|------|---------|

Leishman Stain

Eosin-polychrome Methylene Blue. Romanowsky polychromatic dye. Used to distinguish and identify leukocytes and malaria parasites.

| | |
|------|---------|
| 25 g | LS-P-25 |
|------|---------|

Light Green S.F., C.I. 42095

Acid Green 5, Pencil Green SF, BSC certified stain. For counterstaining according to Papanicolaou and Masson trichrome methods.

| | |
|------|---------|
| 25 g | LG-P-25 |
|------|---------|

Luxol Fast Blue MBSN, C.I. 74180

Luxol Fast Blue, Solvent Blue 38. For use in histology.

| | |
|------|----------|
| 25 g | LFB-P-25 |
|------|----------|

Malachite Green, C.I. 42000

Malachite Green Oxalate, Basic Green 4, Victoria Green, BSC certified stain. Used as counterstain in TB-Stain Cold kit.

| | |
|------|----------|
| 10 g | MAG-P-10 |
| 25 g | MAG-P-25 |

Martius Yellow C.I. 10315

Golden Yellow, Martinsgelb, Acid yellow 24. For staining erythrocytes yellow in trichrome staining methods.

| | |
|------|----------|
| 25 g | MAY-P-25 |
|------|----------|

May-Gruenwald

Polychromatic Romanowsky dye. For blood smears and bone marrow staining.

| | |
|------|---------|
| 25 g | MG-P-25 |
| 50 g | MG-P-50 |

Metanil Yellow, C.I. 13065

Acid yellow 36. For staining cytoplasm, muscle connectors and identifying parietal cells of gastric glands.

| | |
|-------|-----------|
| 25 g | MEY-P-25 |
| 50 g | MEY-P-50 |
| 100 g | MEY-P-100 |

Methyl Green, C.I. 42590

For staining nuclei green.

| | |
|------|----------|
| 25 g | MGR-P-25 |
|------|----------|

Methyl Violet 2B, C.I. 42535

Basic Violet 1, BSC certified stain. For bacterial staining according to Gram.

| | |
|------|-----------|
| 25 g | MV2B-P-25 |
|------|-----------|

Methyl Violet 10B, C.I. 42555

Gentian Violet, Crystal Violet, BSC certified stain. For bacterial staining according to Gram.

| | |
|------|------------|
| 25 g | MV10B-P-25 |
| 50 g | MV10B-P-50 |

Methylene Blue, C.I. 52015

Basic Blue 9, BSC certified stain. Used for preparing Romanowsky polychromatic stain for staining bone marrow and as a counterstain in staining of acid-fast bacteria.

| | |
|-------|----------|
| 25 g | MB-P-25 |
| 100 g | MB-P-100 |

Methylene Green, C.I. 52020

Methylene Green, BSC certified stain. For staining nuclei green.

| | |
|------|----------|
| 25 g | MEG-P-25 |
|------|----------|

Neutral Red, C.I. 50040

Basic Red 5, BSC certified stain. For supravital staining of blood cells using Neutral Red-Janus Green B method.

| | |
|------|---------|
| 25 g | NR-P-25 |
|------|---------|

Nigrosine WS, C.I. 50420

Nigrosine water soluble, Nigrosine B. For use in microbiology and cytology, as negative staining.

| | |
|------|----------|
| 25 g | NGR-P-25 |
|------|----------|

Nuclear Fast Red, C.I. 60760

Kernechtrot. Used for staining nuclei red by creating aluminum complex.

| | |
|-----|--------|
| 5 g | NF-P-5 |
|-----|--------|

Oil Red O, C.I. 26125

Sudan Red 5B, BSC certified stain. For staining lipids.

| | |
|------|----------|
| 25 g | ORO-P-25 |
|------|----------|

Orange II, C.I. 15510

Acid Orange 7, BSC certified stain. Used as a counterstain, as a substitute for Orange G.

| | |
|------|---------|
| 25 g | O2-P-25 |
|------|---------|

Orange G, C.I. 16230

Acid Orange 10, Acid Fast Orange G, BSC certified stain. For staining keratin according to Papanicolaou, counterstain for hematoxylin.

| | |
|-------|----------|
| 25 g | OG-P-25 |
| 100 g | OG-P-100 |

Orcein, C.I. (ed.1) 1242

Natural Red 28, Orcein Synthetic. For staining elastic tissues and viral inclusion bodies of the hepatitis B surface antigen.

| | |
|------|----------|
| 5 g | ORC-P-5 |
| 10 g | ORC-P-10 |

Phloxine B, C.I. 45410

Acid Red 92, Eosin 10B, Eosin Blue, BSC certified stain. For Methylene Blue-Phloxine staining and staining keratin and mucus according to Kreyberg.

25 g PB-P-25

Ponceau S, C.I. 27195

Acid Red 112, Fast Ponceau 2B. For use in electrophoresis.

25 g PS-P-25

Potassium permanganate

Synonym: Potassium (VII) manganate, Condy's crystals. Potassium permanganate for use in microscopy.

50 g KPH-P-50
100 g KPH-P-100

Pyronine B, C.I. 45010

Pyronine B, BSC certified stain. For bacterial, mold and RNA staining.

25 g PYB-P-25

Resorcinol, C.I. 76505

Resorcin, 1,3-benzendiol. Used for differentiation between aldoses and ketoses in Seliwanoff's test by staining ketose sugars dark red.

25 g RES-P-25

Rhodamine B, C.I. 45170

Basic Violet, Acid Red 52. For staining acid-fast bacteria.

25 g RHB-P-25

Rhodanine

DMAB-Rhodanine. For staining deposits of copper bound to proteins.

5 g RHO-P-5

Rose Bengal, C.I. 45440

Acid Red 94, BSC certified stain. For staining in microscopy and histology.

25 g RB-P-25

Safranin O, C.I. 50240

Basic Red 2, Safranin T, Phenazinium, BSC certified stain. Counterstain in the Gram method for detecting Gram-negative bacteria.

25 g SO-P-25

Sirius Red, C.I. 35780

Direct Red 80. Used for staining collagen and amyloid in histology.

25 g SR-P-25

Sudan III, C.I. 26100

Solvent Red 23, Sudan G, Fat Ponceau G, BSC certified stain. For lipid staining according to Lillie-Ashburn.

25 g S3-P-25

Sudan IV, C.I. 26105

Solvent Red 24, Scarlet Red, Biebrich Scarlet R, Fat Ponceau. For staining lipids according to Churukian's modification of Lillie-Ashburn's method.

25 g S4-P-25

Sudan Black B, C.I. 26150

Sudan Black, Solvent Black 3, Fat Black HB. For staining lipids and other hydrophobic materials.

5 g SB-P-5
10 g SB-P-10

Toluidine Blue O, C.I. 52040

Basic Blue 17, Tolonium Chloride, BSC certified stain. For staining mast cells, cartilage and Nissl bodies in the spinal cord.

25 g TB-P-25
100 g TB-P-100

Victoria Blue B, C.I. 44045

Basic Blue 26, Fat Blue B. For use in cytology and histology.

25 g VBB-P-25

Victoria Blue R, C.I. 42563

Basic Blue 8, Fat Blue 4R. For use in microbiology and histology.

25 g VBR-P-25

Water Blue, C.I. 42755

Acid Blue 22, BSC certified stain. For staining connective tissue according to Mallory and Gomori trichrome methods.

25 g WB-P-25

Wright's Stain

Colorants de Wright, BSC certified stain. Part of polychromatic Romanowsky stains commonly used in hematology.

25 g WS-P-25

Xylidine Ponceau, C.I. 16150

Ponceau 2R, Ponceau de Xylidine, BSC certified stain. For Masson trichrome staining.

25 g XP-P-25





Accessories

High quality disposable laboratory equipment including glass, plastic and metal items. The product range includes VitroGnost slides for microscopy, sample transport containers and various other accessories.

VitroGnost® Plus Ultra adhesive microscope slides

These slides are ideal for pathology, cytology, and special immunohistochemistry (IHC). VitroGnost Plus Ultra adhesive slide is a new generation slide with a denser and stronger positive charge on the surface to achieve even better adhesion of the sample. VitroGnost Plus Ultra adhesive slide can be used for many scientific purposes, especially with tissues difficult to fix, like breast tissue and samples of brain and skin. Dimensions: 26x76x1 mm.



72 pcs

103001A72

Lid for cassettes

Cassette embedding stainless steel lid, standard.

| material | packaging | product code |
|----------|-----------|--------------|
| INOX | 1 pc | 51050103 |



Metal histology mold

| dimension | material | packaging | product code |
|-------------|----------|-----------|--------------|
| 7x7x6 mm | INOX | 1 pc | 51050707 |
| 15x15x6 mm | INOX | 1 pc | 51051515 |
| 24x24x6 mm | INOX | 1 pc | 51052424 |
| 30x24x6 mm | INOX | 1 pc | 51053024 |
| 37x24x6 mm | INOX | 1 pc | 51053724 |
| 65x45x15 mm | INOX | 1 pc | BO07-7010 |

Biopsy pads for cassettes, blue

Solvent resistant biopsy pads for small sample processing.

| | |
|----------|---------|
| 1000 pcs | SP-1000 |
|----------|---------|



Polypropylene dissecting boards

Boards for pathohistological tissue dissection made of polypropylene.

| | | |
|---|------|---------|
| Polypropylene, white, dimensions 60x40x1.5 cm | 1 pc | BC001 |
| Polypropylene, dark blue, dimensions 39x29.5x2.5 cm | 1 pc | BC032 |
| Polypropylene, white, dimensions 40x30x1.5 cm | 1 pc | BC061CS |





Bio/Cut microtome blades - high profile

Bio/Cut H35C microtome blades for cryotome, high profile (wide). Ideal for frozen sections. Suitable for both soft and hard tissues.

50 pcs Bio/Cut H35C

Histoblades Duraedge histology blades - high profile

HistoBlades Duraedge blades with Teflon coating - high profile.

50 pcs 7310

Bio/Cut microtome blades - low profile

Bio/Cut microtome blades manufactured in order to provide most precise sectioning.

Bio/Cut 22A microtome blades for ultra-thin sections, kidneys and lymph nodes, narrow blade 50 pcs Bio/Cut 22A

Bio/Cut 35A microtome blades for soft and hard tissues, low profile 50 pcs Bio/Cut 35A

Bio/Cut 35B microtome blades for soft tissues, low profile 50 pcs Bio/Cut 35B

Bio/Cut 35N microtome blades for thin sections, low profile 50 pcs Bio/Cut 35N

Bio/Cut 35R microtome blades for hard tissues, low profile 50 pcs Bio/Cut 35R

Bio/Cut 35S microtome blades for hard tissues, low profile 50 pcs Bio/Cut 35S

Histoblades Duraedge histology blades - low profile

HistoBlades Duraedge blades with Teflon coating - low profile.

50 pcs 7223

Histoblades Encore histology blades - low profile

Extremely sharp new generation Teflon coated blades - low profile.

50 pcs 7380



Microtome oil

Optimal viscosity oil for microtome maintenance.

| | | |
|----------------------|------|--------|
| Microtome oil 50 mL | 1 pc | MO-50 |
| Microtome oil 100 mL | 1 pc | MO-100 |



Polypropylene containers for sample transport

Container with cover for transporting histology samples and organs.

| | | |
|----------|------|---------|
| 500 mL | 1 pc | PK500 |
| 1000 mL | 1 pc | PK1000 |
| 1200 mL | 1 pc | PK1200 |
| 2500 mL | 1 pc | PK2500 |
| 3750 mL | 1 pc | PK3750 |
| 5000 mL | 1 pc | PK5000 |
| 7000 mL | 1 pc | PK7000 |
| 10000 mL | 1 pc | PK10000 |



Contact and ordering information

BioGnost Ltd.,
Medjugorska 59, Zagreb
10040 Croatia (Europe)

WORKING HOURS
Monday through Friday
8:00 to 16:00 (CET)

e-mail: sales@biognost.hr
website: www.biognost.com
phone: +385 1 2409 997
fax: + 385 1 2404 039



Technical support

Technical questions can be answered by emailing our sales representatives at sandra.ban@biognost.hr, pavle.loncar@biognost.hr, viktoria.liber@biognost.hr and andrej.spehar@biognost.hr.

They will provide you with any MSDS or product information required, also available on our website at www.biognost.com.

Frequently Asked Questions (FAQ) section is available on our website.



Shipping information

Shipping and handling charges are prepaid and added to the invoice. If requested, we can use your account number or carrier of your choice, and also ship on your third-party freight accounts (Incoterms 2020). Charges vary according to destination, weight and content, and are available upon request.

Our employees are trained to classify, package, mark and label products identified as dangerous goods according to ADR, IMDG and IATA to ensure safe transport.



Index

| | |
|-------------------------------------|----|
| 4% Formaldehyde | 8 |
| 4% Formaldehyde, alcoholic solution | 51 |
| 4% NB Formaldehyde | 8 |
| 10% Formaldehyde | 9 |
| 10% NB Formaldehyde | 9 |

A

| | |
|--|----|
| Acetic acid, 0.5% solution | 53 |
| Acetic acid, 1% solution | 53 |
| Acetic acid, Colloidal Iron (A) solution | 53 |
| Acetic acid, Colloidal Iron (B) solution | 53 |
| Acetic buffer, Grimelius | 51 |
| Acid alcohol | 24 |
| Acid alcohols | 51 |
| Acid solution | 51 |
| Acridine Orange, C.I. 46005 | 58 |
| Acridine Orange solution | 51 |
| A.F.O.G. kit | 41 |
| A.F.O.G. reagent | 51 |
| Alcian Blue 8 GX, C.I. 74240 | 58 |
| Alcian Blue - P.A.S. kit | 41 |
| Alcian Blue pH 1.0 kit | 41 |
| Alcian Blue pH 2.5 kit | 41 |
| Alcian Blue solution 1% | 51 |
| Alcian Blue solution pH 1.0 | 51 |
| Alcian Blue solution pH 2.5 | 51 |
| Alcian Green | 58 |
| Alcian Yellow, C.I. 12840 | 58 |
| Alcian Yellow Toluidine Blue kit | 41 |
| Alcoholic Eosin Y solutions | 25 |
| Alkaline alcohol, solution | 51 |
| Amido Black 10B, C.I. 20470 | 58 |
| Ammonium iron sulfate, solution | 52 |
| Ammonium water | 52 |
| Aniline, alcoholic solution | 52 |
| Aniline Blue, C.I. 42780 | 58 |
| Aniline Blue reagent | 52 |
| Aniline Blue, solution | 52 |
| Auramine O, C.I. 41000 | 58 |
| Azan reagent | 52 |
| Azan Trichrome kit | 41 |
| Azocarmine G, C.I. 50085 | 58 |
| Azocarmine, solution | 52 |
| Azophloxine, C.I. 18050 | 58 |
| Azure A, C.I. 52005 | 58 |
| Azure B, C.I. 52010 | 58 |
| Azure C, C.I. 52002 | 58 |
| Azure II | 58 |
| Azure II Eosinate | 59 |

B

| | |
|---|--------|
| Basic Activation buffer | 52 |
| Benedict's reagent | 52 |
| Biebrich Scarlet-Acid Fuchsin reagent | 52 |
| Biebrich Scarlet, C.I. 26905 | 59 |
| Bielschowsky kit | 42 |
| BioClear | 13 |
| BioClear New | 13 |
| Bio/Cut microtome blades - high profile | 64 |
| Bio/Cut microtome blades - low profile | 64 |
| Bio-Diff 1 reagent | 28 |
| Bio-Diff 1 RTU reagent | 29 |
| Bio-Diff 2 reagent | 28 |
| Bio-Diff 2 RTU reagent | 29 |
| Bio-Diff 3 reagent | 28 |
| Bio-Diff 3 RTU reagent | 29 |
| Bio-Diff kit | 28 |
| Bio-Diff RTU kit | 29, 42 |
| BioFix | 9 |

| | |
|---------------------------------------|--------|
| BioFix C | 9 |
| BioFix GL | 9 |
| BioFix HD | 9 |
| BioForm S | 9 |
| BioGram 4 kit | 34 |
| BioGram ECO kit | 35 |
| BioGram Histo kit | 42 |
| BioMount | 18 |
| BioMount Aqua | 19, 33 |
| BioMount C | 19 |
| BioMount DPX | 17 |
| BioMount DPX High | 17 |
| BioMount DPX Low | 17 |
| BioMount DPX New | 17 |
| BioMount High | 18 |
| BioMount M | 18 |
| BioMount New | 18 |
| BioMount New Low | 18 |
| Biopsy pads for cassettes, blue | 63 |
| BioSchiff reagent | 52 |
| BioWax 52/54 | 14 |
| BioWax 56/58 | 14 |
| BioWax Blue | 14 |
| BioWax Plus 56/58 | 14 |
| Bismarck Brown Y, C.I. 21000 | 59 |
| Bluing reagent | 24 |
| Borax, solution | 52 |
| Bouin's solution | 10 |
| Brilliant Cresyl Blue, C.I. 51010 | 59 |
| Brilliant Cresyl Blue, solution | 31 |
| Brilliant Green Certified, C.I. 42040 | 59 |
| Buffer solution, pH 6.4 | 30 |
| Buffer solution, pH 6.8 | 30 |
| Buffer solution, pH 7.2 | 29 |
| Buffer tablets pH 6.8 | 30 |
| Buffer tablets pH 7.2 | 29 |

C

| | |
|--|----|
| Carbol Fuchsin Gram 10x concentrated | 35 |
| Carmine, C.I. 75470 | 59 |
| Carminic acid, C.I. 75470 | 59 |
| Celestine Blue, C.I. 51050 | 59 |
| Chromotrope 2R / Aniline Blue reagent | 53 |
| Chromotrope 2R, C.I. 16570 | 59 |
| Chromotrope 2R / Fast Green F.C.F. reagent | 53 |
| CitoFix | 10 |
| CitoSpray | 10 |
| CitoSpray solution | 10 |
| Clean Lens, solution 1 | 21 |
| Clean Lens, solution 2 | 21 |
| Clean Lens, solution 3 | 21 |
| Colloidal Iron kit | 42 |
| Colloidal Iron, stock solution | 53 |
| Congo Red, C.I. 22120 | 59 |
| Congo Red Highman kit | 42 |
| Congo Red Puchtler kit | 42 |
| Congo Red Puchtler reagent | 53 |
| Congo Red reagent | 53 |
| Cresyl Violet Acetate Certified | 59 |
| Cresyl Violet, solution | 53 |
| CryoFix Gel | 15 |
| Cryo Spray | 15 |
| Crystal Violet, C.I. 42555 | 59 |
| CytoGnost kit | 31 |

D

| | |
|---|----|
| DeHematin | 9 |
| DeMold solution | 15 |
| DeMold Spray | 15 |
| Developer stock solution | 53 |
| Differentiation reagents based on acetic acid | 53 |
| Differentiation reagents based on hydrochloric acid | 53 |
| Duoline marker pen | 7 |

| | |
|--|--------|
| E | |
| EA 31 reagent, Pap 3A | 27 |
| EA 50 reagent, Pap 3B | 27 |
| EA 65 reagent, Pap 3C | 27 |
| EA 65 reagent, Pap 3D | 27 |
| Ehrlich aldehyde reagent | 53 |
| Elastica-Van Gieson kit | 43 |
| Eosin and nigrosine vital kit | 43 |
| Eosin B, C.I. 45400 | 59 |
| Eosin Contrast | 25 |
| Eosin Contrast PLUS | 25 |
| Eosin-nigrosin vital | 53 |
| Eosin Y aqueous solutions | 25 |
| Eosin Y, C.I. 45380 | 59 |
| Eosin Y instant | 25 |
| Eosin Y, isotonic solution | 53 |
| Eriochrome Cyanine R, C.I. 43820 | 59 |
| ErioGnost reagent | 24, 53 |
| F | |
| Fast Green F.C.F., C.I. 42053 | 59 |
| Fast Green F.C.F. contrast reagent | 53 |
| Fast Green F.C.F. reagent | 53 |
| Ferri reagents, Verhoeff B | 53 |
| Ferri reagent, Weigert B | 53 |
| Feulgen kit | 43 |
| Field kit | 31, 43 |
| Field, solution A | 53 |
| Field, solution B | 53 |
| Formaldehyde 37% for use in histology | 9 |
| Fouchet-Van Gieson kit | 43 |
| Fuchsin Acid, C.I. 42685 | 59 |
| Fuchsin Acid reagent | 53 |
| Fuchsin Acid Van Gieson reagent | 54 |
| Fuchsin Basic, C.I. 42510 | 59 |
| Fuchsin Basic reagent | 54 |
| Fuchsin Basic - Schiff, C.I. 42500 | 59 |
| Fuchsin Basic, solution | 54 |
| G | |
| Gabbet reagent | 37 |
| Gelatin, gel in transport jar | 54 |
| Gentian Violet, C.I. 42555 | 59 |
| Giemsa | 59 |
| Giemsa HP kit | 43 |
| Giemsa solution | 30 |
| Glutaraldehyde, 4% buffered solution | 54 |
| Gold chloride solutions | 54 |
| Gomori Trichrome kit | 44 |
| Gram Crystal Violet 1% solution | 34 |
| Gram Crystal Violet, phenol-free reagent | 35 |
| Gram Decolorizer solution 1 | 35 |
| Gram Decolorizer solution 2 | 34 |
| Gram Decolorizer solution 3 | 35 |
| Gram Lugol solution | 35 |
| Gram Lugol solution, stabilized | 34 |
| Gram Safranin 5x concentrated | 35 |
| Gram Safranin solution | 34 |
| Gram sodium hydrogencarbonate, solution | 35 |
| Grimelius kit | 44 |
| H | |
| Hayem's solution | 31 |
| H.B.F.P. kit | 44 |
| HCL Pap reagent | 24 |
| HCL reagent, A.F.O.G. | 53 |
| HCL reagent, Alcian Blue | 53 |
| HCL reagent, Colloidal Iron | 53 |
| HCL reagent, Feulgen | 53 |
| HCL reagent, HemoGnost Perls | 53 |
| HCL reagent, P.A.S. | 53 |
| HCL reagent, Weigert-Van Gieson | 53 |
| Hematoxylin CAR | 24 |
| Hematoxylin, C.I. 75290 | 60 |
| Hematoxylin COL | 24 |
| Hematoxylin G1 | 23, 27 |
| Hematoxylin G2 | 23, 27 |
| Hematoxylin G3 | 23, 27 |
| Hematoxylin H | 23 |
| Hematoxylin HP, Pap 1A | 26 |
| Hematoxylin HP, Pap 1B | 26 |
| Hematoxylin Instant kit | 24 |
| Hematoxylin M | 23 |
| Hematoxylin ML | 23 |
| Hematoxylin P.T.A. | 24 |
| Hematoxylin P.T.A. kit | 44 |
| Hematoxylin, Verhoeff A | 54 |
| Hematoxylin, Weigert A | 54 |
| Hematoxylin W kit | 24, 44 |
| HemoGnost Perls kit | 45 |
| Histanol 50 | 12 |
| Histanol 70 | 12 |
| Histanol 80 | 12 |
| Histanol 95 | 12 |
| Histanol 96 | 12 |
| Histanol 100 | 12 |
| Histanol EP 95 | 12 |
| Histanol EP 100 | 12 |
| Histanol G | 12 |
| Histanol IP | 12 |
| Histanol M | 12 |
| Histoblades Duraedge histology blades - high profile | 64 |
| Histoblades Duraedge histology blades - low profile | 64 |
| Histoblades Encore histology blades - low profile | 64 |
| Hydroquinone-sodium sulphite | 54 |
| Hydroquinone, solution | 54 |
| I | |
| Immersion oil | 20 |
| Immersion oil, tropical grade | 21 |
| Immersion oil type 37 | 21 |
| Immersion oil type A | 20 |
| Immersion oil type C | 21 |
| Immersion oil type FF | 21 |
| ImmunoForm | 9 |
| Iodine solution, Verhoeff C | 54 |
| Iron chloride, solution | 54 |
| J | |
| Janus Green B, C.I. 11050 | 60 |
| Jenner Stain | 60 |
| K | |
| Kit for rapid HE staining of frozen and paraffin-embedded sections | 24 |
| Kovac indole reagent | 54 |
| L | |
| Lactophenol Blue solution | 39 |
| Lactophenol solution | 39 |
| Ladewig's reagent | 54 |
| Ladewig Trichrome kit | 45 |
| Leishman's solution | 31 |
| Leishman Stain | 60 |
| LeukoGnost ACP | 32 |
| LeukoGnost ALP | 32 |
| LeukoGnost Fixative | 33 |
| LeukoGnost Fixative ECO | 33 |
| LeukoGnost Fixative GL | 33 |
| LeukoGnost HEM | 33 |
| LeukoGnost MPO | 32 |
| LeukoGnost NSE | 32 |

| | | | |
|--|----|---|--------|
| LeukoGnost PAS | 32 | Orange G, C.I. 16230 | 60 |
| LeukoGnost PERLS | 32 | Orange II, C.I. 15510 | 60 |
| LeukoGnost PLUS | 33 | Orange II reagent, Pap 2B | 26 |
| LeukoGnost SPE | 32 | Orcein, C.I. (ed.1) 1242 | 60 |
| LeukoGnost SPE kit | 45 | Orcein kit | 47 |
| LeukoGnost SPENSE | 32 | Orcein reagent | 55 |
| Lid for cassettes | 62 | OsteoFast 1 | 11 |
| Light Green S.F., C.I. 42095 | 60 | OsteoFast 2 | 11 |
| Light Green S.F. reagent | 54 | OsteoSens | 11 |
| Liquid Blocker marker pen | 7 | Oxalic acid solutions | 55 |
| Lithium carbonate, Luxol | 54 | | |
| Lithium carbonate, solution | 54 | P | |
| Lugol's C solution | 35 | Paraldehyde Fuchsin kit | 47 |
| Lugol's solution, acid | 35 | Paraldehyde Fuchsin reagent | 55 |
| Lugol's solution, M.I.F. | 54 | P.A.S. Diastase kit | 40, 47 |
| Lugol's solution | 35 | P.A.S. kit | 47 |
| Luxol Fast Blue kit | 45 | Periodic acid, solution | 55 |
| Luxol Fast Blue MBSN, C.I. 74180 | 60 | Perls-Van Gieson kit | 48 |
| Luxol Fast Blue, solution | 54 | Phloxine B, C.I. 45410 | 61 |
| | | Phosphomolybdic acid, 1% solution | 55 |
| M | | Phosphomolybdic acid, 5% solution | 55 |
| Malachite Green, C.I. 42000 | 60 | Phosphotungstic acid, 1% solution | 55 |
| Mallory Trichrome kit | 45 | Phosphotungstic acid, 5% solution | 55 |
| Martius Scarlet Blue (MSB) kit | 46 | Picric acid in acetone, solution | 55 |
| Martius Yellow C.I. 10315 | 60 | Picro-Sirius Red kit | 48 |
| Martius Yellow, solution | 54 | Picro-Sirius Red reagent | 55 |
| Masson Fontana kit | 46 | Polypropylene containers for transporting samples | 65 |
| Masson-Goldner Trichrome kit | 46 | Polypropylene dissecting boards | 63 |
| Masson Trichrome kit | 46 | Ponceau S, C.I. 27195 | 61 |
| Mayer's mucicarmine | 54 | Ponceau S, solution | 55 |
| May-Gruenwald | 60 | Potassium hexacyanoferrate, Colloidal Iron solution | 55 |
| May-Gruenwald solution | 30 | Potassium hexacyanoferrate, solution | 56 |
| Metal histology mold | 63 | Potassium hydroxide, solution | 56 |
| Metanil Yellow, C.I. 13065 | 60 | Potassium hydroxide, solution for fungi | 39 |
| Metanil Yellow, solution | 54 | Potassium permanganate | 61 |
| Methenamine, solution | 55 | Potassium permanganate, solutions | 56 |
| Methyl Blue Eosin, solution | 55 | Propylene glycol | 56 |
| Methylene Blue 10X concentrated | 36 | Propylene glycol, 85% | 56 |
| Methylene Blue, C.I. 52015 | 60 | P.T.A. - P.M.A. reagent | 56 |
| Methylene Blue Loeffler's reagent | 36 | Pyronine B, C.I. 45010 | 61 |
| Methylene Green, C.I. 52020 | 60 | | |
| Methyl Green, C.I. 42590 | 60 | R | |
| Methyl Violet 2B, C.I. 42535 | 60 | Rapid Pap staining kit | 27 |
| Methyl Violet 10B, C.I. 42555 | 60 | Reagent for differentiation in Gomori Trichrome kit | 53 |
| M.F. reagent | 54 | Reagent for differentiation in Verhoeff kit | 56 |
| Micro Clean | 14 | Reducing buffer, solution | 56 |
| Microtome oil | 65 | Resorcine Fuchsin reagent | 56 |
| M.I.F. kit | 45 | Resorcinol, C.I. 76505 | 61 |
| Movat kit | 46 | Reticulin Contrast kit | 48 |
| Mucicarmine kit | 46 | Reticulin kit | 48 |
| Myco-Mount Blue | 39 | Rhodamine B, C.I. 45170 | 61 |
| Myco-Mount Red | 39 | Rhodanine | 61 |
| | | Rhodanine kit | 48 |
| | | Rhodanine reagent | 56 |
| N | | Rose Bengal, C.I. 45440 | 61 |
| NB Formaldehyde concentrated 5x | 9 | | |
| Neutral Red, C.I. 50040 | 60 | S | |
| Neutral Red reagent | 55 | Saccommanno's fixative | 10 |
| Nigrosine, isotonic solution | 55 | Safranin O, C.I. 50240 | 61 |
| Nigrosine WS, C.I. 50420 | 60 | Safranin O kit | 48 |
| Nuclear Fast Red, C.I. 60760 | 60 | Safranin O reagent | 56 |
| Nuclear Fast Red (Kernechtrot) reagent | 55 | Scott's solution | 24 |
| | | Shorr's solution | 27 |
| O | | Silver nitrate solutions | 56 |
| OG-6 reagent, Pap 2A | 26 | Sirius Red, C.I. 35780 | 61 |
| OG-EA31 Pap reagent | 27 | Sodium acetate, solution | 56 |
| OG-EA50 Pap reagent | 27 | Sodium hydroxide, solution | 56 |
| Oil Red O, C.I. 26125 | 60 | Sodium metabisulfite solutions | 56 |
| Oil Red O kit | 47 | Sodium tetraborate, solution | 56 |
| Oil Red O reagent | 55 | Sodium thiosulfate solution | 56 |
| Orange G, 1% solution | 55 | Sperm-Diff RTU kit | 49 |
| Orange G / Aniline Blue reagent | 55 | | |

| | |
|------------------------------------|----|
| Sudan Black B, C.I. 26150 | 61 |
| Sudan Black B ECO | 32 |
| Sudan Black B kit | 32 |
| Sudan Black B lipid kit | 49 |
| Sudan Black B reagent | 57 |
| Sudan Black B, solution | 56 |
| Sudan III, C.I. 26100 | 61 |
| Sudan IV, C.I. 26105 | 61 |
| Sulfuric acid solutions | 57 |
| Sulphosalicylic acid, 20% solution | 57 |

T

| | |
|-----------------------------------|--------|
| TB Armand reagent | 37 |
| TB Auramine O reagent | 38 |
| TB Auramine-Rhodamine reagent | 38 |
| TB Carbol Fuchsin reagent | 36, 37 |
| TB Decolorizer | 36, 37 |
| TB Decolorizer Fluorescent | 38 |
| TB-Fuchsin reagent | 36 |
| TB Malachite Green reagent | 37 |
| TB Permanganate reagent | 38 |
| TB-Stain Auramine O kit | 38 |
| TB-Stain Cold kit | 37 |
| TB-Stain ECO kit | 36 |
| TB-Stain Fluorescent kit | 38 |
| TB-Stain Histo kit | 49 |
| TB-Stain Hot kit | 36 |
| TB-Stain Quick kit | 37 |
| ThromboGnost solution | 31 |
| Tissue marking dyes | 7 |
| Tissue marking dyes, 5 colour set | 7 |
| Tissue marking dyes, 7 colour set | 7 |
| Tissue Soft | 15 |
| Toluidine Blue O, C.I. 52040 | 61 |
| Toluidine Blue, solution | 57 |
| Trichloroacetic acid, solution | 57 |
| Tuerk's solution | 31 |

U

| | |
|---------------------|----|
| UriGnost S kit | 49 |
| UriGnost SM kit | 49 |
| UriGnost SM reagent | 57 |
| UriGnost S reagent | 57 |

V

| | |
|---|----|
| Van Gieson Trichrome kit | 49 |
| Verhoeff kit | 50 |
| Victoria Blue B, C.I. 44045 | 61 |
| Victoria Blue R, C.I. 42563 | 61 |
| VitroGnost® Plus Ultra adhesive microscope slides | 62 |
| Von Kossa kit | 50 |

W

| | |
|--------------------------------------|----|
| Warthin Starry kit | 50 |
| Water Blue, C.I. 42755 | 61 |
| Weigert-Van Gieson kit | 50 |
| Wright-Giemsa solution | 31 |
| Wright-Giemsa, solution for reptiles | 31 |
| Wright's solution | 31 |
| Wright's Stain | 61 |

X

| | |
|------------------------------|----|
| Xylidine Ponceau, C.I. 16150 | 61 |
|------------------------------|----|

Be a part of
our global network
of satisfied customers

BioGnost Ltd.

For development, manufacture and
marketing of pharmaceuticals

Medjugorska 59, 10040 Zagreb, Croatia, EU
phone: +385 1 2409 997, fax: +385 1 2404 039
sales@biognost.hr • www.biognost.com

