

CHIRASORB is a mid-term absorbable, coated, synthetic, multifilament suture made of 90% glycolide and 10% L-lactide. The substances contained in the coating and suture are nonantigenic and noncollagenous.

Sterile Chirasorb is available in violet or natural. The product elicits a minimal acute inflammatory reaction in tissues.

Progressive loss of tensile strength and absorption of Chirasorb occurs by hydrolysis, when the polymer is degraded to glycolic and lactic acids, which are subsequently absorbed and metabolised by the body.



Indication:	General surgery, Gynaecology, Ophthalmic, Skin Closure (e.g. intracutan- subcutaneous), Muscular layer and Ligation.	eous,				
Contraindication:	Chirasorb should not be used in tissues subject to expansion, stretching or distension, or which might require long-term mechanical support.					
Sizes:	10/0 to 6 USP					
Structure:	Multifilament					
Absorption method:	Hydrolysis					
Strength Retention Profile:	MID-TERM HEALING (35 days) 1 Day 100% 14 Days 75%					
Mass Absorption profile:	21 Days 50%					

*

下名用

UK03

CTI group

CHIRASORB

biger BROBARIKS

3	Needle	EP	USP	Length	Colour	Cat. No	Other information
	HR: ½ circle, ro	und body needle					
)		4	1	1 x 45 cm	Violet	LV 0177	1.57
2				- 11			
2		5	2	5 x 45 cm	Violet	LV 2250	
1							
5				and the set			
٤	HR 40	2	2/0	1 x 75 cm	Violet	LV 0275	
2	\ • /	3 3.5	2/0 0	1 x 75 cm	Violet	LV 0276	
	\bigvee	4	1	1 x 75 cm	Violet	LV 0277	
ć			2/0	1 x 90 cm	Violet	LV 1475	
		≥ 3 3.5	0	1 x 90 cm	Violet	LV 1476	
		4	1	1 x 90 cm	Violet	LV 1477	
		5	2	1 x 90 cm	Violet	LV 1478	
			2/0	1 x 120 cm	Violet	LV 1479	
		3	210		VIOLO		
	HR 43	п					
	(°	5	2	1 x 75 cm	Violet	LV 0479	
					C.		6
	110.40	4	1	1 x 75 cm 1 x 75 cm	Violet Violet	LV 1396 LV 1400	
	HR 48	5 3.5 Л	2 0	1 x 75 cm	Violet	LV 1402	
		3.5	0	1 x 90 cm	Violet	LV 1403	8.7
		4 5	1	1 x 90 cm 1 x 90 cm	Violet	LV 1401 LV 1399	
		5	2	1 x 90 cm	Violet	LV 1399	
	HR 48 A						
) 4	1	4 x 45 cm	Violet	LV 0793	Take-off needl
							• 11 a)