

Designed for above the knee interventions

BioPath™ 035 offers excellent pushability, trackability and crossability due to a low balloon profile, low tip entry profile and a hydrophilic coating on the distal shaft of the catheter.

BioPath 035, with its unique shellac coating, delivers a designated paclitaxel dosage where it matters most – across the target lesion.

BioPath 035 is indicated for:

- > De-novo lesions
- > Restenosis after realisation of balloon and /or stent PTA
- > Pre- and post-dilatation in case of peripheral stent implantation

BIOPATH™ 035
PACLITAXEL ELUTING PTA BALLOON CATHETER (OTW)

Latest generation paclitaxel-eluting balloon
for peripheral interventions

Ordering Information

Balloon diameter (mm)	Balloon length (mm)						
	20	40	60	80	100	120	150
4.0	BPTH-35-4020 S	BPTH-35-4040 S	BPTH-35-4060 S	BPTH-35-4080 S	BPTH-35-40100 S	BPTH-35-40120 S	BPTH-35-40150 S
	BPTH-35-4020 L	BPTH-35-4040 L	BPTH-35-4060 L	BPTH-35-4080 L	BPTH-35-40100 L	BPTH-35-40120 L	BPTH-35-40150 L
5.0	BPTH-35-5020 S	BPTH-35-5040 S	BPTH-35-5060 S	BPTH-35-5080 S	BPTH-35-50100 S	BPTH-35-50120 S	BPTH-35-50150 S
	BPTH-35-5020 L	BPTH-35-5040 L	BPTH-35-5060 L	BPTH-35-5080 L	BPTH-35-50100 L	BPTH-35-50120 L	BPTH-35-50150 L
6.0	BPTH-35-6020 S	BPTH-35-6040 S	BPTH-35-6060 S	BPTH-35-6080 S	BPTH-35-60100 S	BPTH-35-60120 S	BPTH-35-60150 S
	BPTH-35-6020 L	BPTH-35-6040 L	BPTH-35-6060 L	BPTH-35-6080 L	BPTH-35-60100 L	BPTH-35-60120 L	BPTH-35-60150 L
7.0	BPTH-35-7020 S	BPTH-35-7040 S	BPTH-35-7060 S	BPTH-35-7080 S	BPTH-35-70100 S	BPTH-35-70120 S	BPTH-35-70150 S
	BPTH-35-7020 L	BPTH-35-7040 L	BPTH-35-7060 L	BPTH-35-7080 L	BPTH-35-70100 L	BPTH-35-70120 L	BPTH-35-70150 L
8.0	BPTH-35-8020 S	BPTH-35-8040 S	BPTH-35-8060 S	BPTH-35-8080 S	BPTH-35-80100 S	S: 80 cm usable catheter length L: 135 cm usable catheter length	
	BPTH-35-8020 L	BPTH-35-8040 L	BPTH-35-8060 L	BPTH-35-8080 L	BPTH-35-80100 L		

1. Axel D.L. et al. Circulation 1997; 96:636-45
2. A. Posa et al. Catheterization and Cardiovascular Interventions 76:395-403 (2010)
3. Data on file

BioPath™ 035 paclitaxel-eluting balloon catheter is CE Mark approved.

CAUTION: The law restricts these devices to sale by or on the order of a physician. Indications, contraindications, warnings and instructions for use can be found in the product labeling supplied with each device.

BioPath is a trademark or registered trademark of Biosensors International Group, Ltd. All cited trademarks are the property of their respective owners. Les informations relatives à ce produit ne sont pas destinées aux professionnels de santé exerçant en France - Any information related to this product is not intended for health care professionals who practice in France.

Not available for sale in the United States and certain other countries.
© 2014 Biosensors International Group, Ltd. All rights reserved.

www.biosensors.com

BIOSENSORS
INTERNATIONAL™

BIOSENSORS EUROPE SA
Rue de Lausanne 29
1110 Morges
Switzerland
Tel: +41 (0)21 804 80 00
Fax: +41 (0)21 804 80 01

**BIOSENSORS INTERVENTIONAL
TECHNOLOGIES PTE LTD**
36 Jalan Tukang
Singapore 619 266
Tel: +65 6213 5777
Fax: +65 6213 5737

11275000N Rev.02

BIOPATH™ 035
PACLITAXEL ELUTING PTA BALLOON CATHETER (OTW)

Latest generation paclitaxel-eluting balloon
for peripheral interventions

the right
reach

BIOSENSORS
INTERNATIONAL™

The right reach

The treatment process

With balloon dilatation, the injuries to the arterial wall initiate an inflammatory reaction with an excretion of growth factors which trigger the onset of cell division and smooth muscle cell migration.

Paclitaxel prevents restenosis by stabilizing microtubular formation and thus prevents the cells going through the phases of replication, resulting in the inhibition of cell division.

Paclitaxel reduces the excretion of the platelet derived growth factor (PDGF) that mediates vascular smooth muscle cell migration to the intima¹.


Hydrophilic shaft coating

The BioPath™ 035 balloon coating

- ▶ The BioPath 035 balloon coating consists of a 1:1 mixture of paclitaxel (3 µg/mm²) and shellac, a natural resin approved by the FDA (GRAS), and by Europe (E904) as a food additive.
- ▶ BioPath 035 delivers the designated concentration of paclitaxel locally to the arterial tissue.
- ▶ The properties of shellac protect the paclitaxel during transition and placement.

Delivering the drug


Once in contact with blood, the shellac and paclitaxel coating swells and begins to open, facilitating the pressure-induced transfer of the paclitaxel.



Delivering the paclitaxel drug


Immediately after the controlled PTA, injury to the vessel wall key pathways contribute to the formation of neointimal hyperplasia. Injury from cracking plaque can lead to narrowing of the lumen. The paclitaxel dose will act over the short term to inhibit cell re-growth.

- 1 Pre-dilatation prepares the way for the delivery of paclitaxel from the BioPath 035 balloon surface.
- 2 BioPath 035 is advanced to the lesion site.
- 3 Once the operator is satisfied with the position of BioPath 035 across the lesion, an inflation at 6 bar for at least 60 seconds will deliver the paclitaxel through the cracked plaque and onto the vessel wall.
- 4 BioPath 035 is then withdrawn. The shellac carrier remains on the balloon surface.




Designed for above the knee interventions

Over-the-wire (OTW) catheter



Low Tip Entry Profile

Coronary² and Peripheral³ – Tissue paclitaxel levels 45 minutes post-dilatation



A choice of balloon length and diameter, on two catheter shaft lengths

Available balloon diameters	4.0, 5.0, 6.0, 7.0 and 8 mm
Balloon lengths	20, 40, 60, 80, 100, 120 and 150 mm
Usable catheter lengths	S: 80 cm or L: 135 cm
Recommended guide wire	0.035"

For illustration purpose only - not to scale.