F CHEMI-PHARM®

CHEMISEPT MED

Alcoholic hand disinfectant

Chemisept Med is rapid ethanol-based hand disinfectant for hygienic and surgical hand disinfection. It is developed for professional use and is suitable for any situations with high hygiene requirements. It does not contain colourants and fragrance, which makes it especially suitable for people with sensitive skin. **Chemisept Med** is an effective solution for quick disinfection of the hands.

Chemisept Med has a broad spectrum of bactericidal (incl. MRSA, VRE), mycobactericidal (incl. Mycobacterium avium and Mycobacterium terrae), fully virucidal (incl. all enveloped viruses, Adeno, Noro, Rota and Polio) and yeasticidal activity.



Chemisept Med has been dermatologically tested to be suitable for both normal and sensitive skin. It contains emollients for skin-care, such as glycerine and panthenol, which help rapidly restored the normal microflora, moisture balance and protective functions of the skin

APPLICATION AREAS:

Chemisept Med is alcohol-based hand rub for hygienic and surgical hand disinfection. Perfect product in higher hygiene requirements, where hand antisepsis is the most important preventive measure to avoid infection.

Main areas of usage:

- Hospitals and other medical institutions
- Operating theatres and procedure rooms
- Intensive care unit
- In accident and emergency departments
- Dialysis departments
- Family doctor offices, first-aid stations
- Ambulances
- Laboratories and diagnostics.
- Dentist's offices
- For home care of patient
- Food industry and food processing areas
- Pharmaceutical and cosmetics industry

USAGE:

Chemisept Med is rubbed undiluted onto dry hands. Hands must be completely covered during the application time. Keep special attention to fingertips and thumbs.

Hygienic hand disinfection acc. to EN 1500:

Apply 3 ml of **Chemisept Med** onto previously washed and dried hands and rub until the hands are dry. Contact time is 30 seconds.

Surgical hand disinfection acc. to EN 12791:

Rub at least 3 ml of **Chemisept Med** onto hands and forearms twice. (It is important that the entire area to be disinfected is covered with the solution).

Contact time is 1,5 minutes (45 seconds twice). Both hands and forearms should be disinfected.

MICROBIOLOGY EFFICACY:

| Efficacy | Test method | Contact time |
|--------------------------------|-------------|--------------|
| Hygienic hand disinfection | EN1500 | 30 sec |
| Surgical hand disinfection | EN12791 | 90 sec |
| Bactericidal (incl. MRSA, VRE) | EN13727 | 15 sec |
| Yeasticidal | EN13624 | 15 sec |
| Mycobactericidal | EN14348 | 20 sec |
| All enveloped viruses | EN14476 | 15 sec |
| Norovirus, Rotavirus | EN14476 | 15 sec |
| Adenovirus | EN14476 | 30 sec |
| Poliovirus | EN14476 | 60 sec |
| Fully virucidal | EN14476 | 60 sec |

INGREDIENS:

100 g of product contains active ingredients: ethanol 72,5 g, isopropanol 7,5 g. **pH 6.5 - 7.5**

PRECAUTIONS:

Highly flammable liquid and vapour. Keep out of reach of children. Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Keep container tightly closed. Store in a well-ventilated place. Keep cool. Dispose of contents and / or containers in accordance with regulations on hazardous waste or packaging and packaging waste respectively

PACKAGING:

