



## Tryptose Broth

M177

Tryptose Broth is recommended for the cultivation primarily of *Brucella*.

### Composition\*\*

Ingredients	Gms / Litre
Tryptose	20.000
Dextrose	1.000
Sodium chloride	5.000
Final pH ( at 25°C)	7.3±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 26 grams in 1000 ml distilled water. If desired, add 0.5 - 1% agar to the medium. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

### Principle And Interpretation

Huddleson used Tryptose media for the isolation of *Brucella* species from man (1). Tryptose containing media, rather than the conventionally used meat infusion media have been used for the enumeration and isolation of *Brucella* species (2, 3).

Tryptose Broth is also recommended by APHA (4) and FDA (5). This medium can be used as general purpose media for cultivation of wide variety of organisms. It can also be supplemented with defibrinated blood (sheep, horse) to prepare blood containing medium for the isolation of fastidious organisms like *Brucella*. Tryptose Broth can be supplemented with 0.1% agar for the cultivation of anaerobes.

Dextrose is the source of energy. Tryptose serves as nitrogen source while sodium chloride maintains osmotic equilibrium.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Colour and Clarity of prepared medium

Basal Medium : Yellow coloured, clear solution. With addition of 5% v/v sterile defibrinated blood, cherry red coloured, opaque solution forms in tubes.

#### Reaction

Reaction of 2.6% w/v aqueous solution at 25°C. pH : 7.3±0.2

#### pH

7.10-7.50

#### Cultural Response

M177: Cultural characteristics observed after an incubation at 35-37°C for 48-72 hours with added 5% v/v sterile defibrinated blood in presence of 10% Carbon dioxide (CO<sub>2</sub>).

Organism	Inoculum (CFU)	Growth
<b>Cultural Response</b>		
<i>Brucella melitensis</i> ATCC 4309	50-100	good-luxuriant
<i>Brucella suis</i> ATCC 4314	50-100	good-luxuriant
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	good-luxuriant
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good-luxuriant

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Huddleson I. F., 1943, Brucellosis in man and animals, rev., Ed., The Commonwealth Fund, New York, N.Y.
2. Ruiz Castañeda M., 1947, Proc. Soc. Exp. Biol. Med., 64:114.
3. Huddleson I. F., 1939, Brucellosis in Man and Animals, Oxford University Press, Oxford, England.
4. Downes F. P. and Ito K., (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods. 4th Ed. American Public Health Association, Washington, D.C.
5. U.S. Food and Drug Administration, 1995, Bacteriological Analytical Manual, 8th Ed., AOAC International, Gaithersburg, Md.

Revision : 1 / 2011



### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.