

PHENYLALANINE AGAR

Medium for enterobacteria differentiation.

TYPICAL FORMULA (g/l)

Yeast Extract	3.0
Disodium Phosphate	1.0
Sodium Chloride	5.0
DL-Phenylalanine	2.0
Agar	15.0
Final pH = 7.3 ± 0.2 at 25 °C.	

DIRECTIONS

Suspend 26.0 g of powder in 1 liter of distilled or deionized water. Heat until completely dissolved. Distribute into final tubes. Sterilize in autoclave at 121°C for 15 minutes. Allow the medium to solidify in a slanted position.

DESCRIPTION

PHENYLALANINE AGAR is a medium recommended for the differentiation of members of the *Proteus* and *Providencia* groups from other enterobacteria.

TECHNIQUE

Inoculate the slant with test organisms and incubate at 36 ± 1°C for 18-24 hours. Add 3-5 drops of Ferric Chloride 10% (code 80272) and 3-5 drops of a 0.1 N HCl solution to a 24 hours culture. Rotate the tubes to wet and loosen the growth.

A positive test is indicated by the formation of a characteristic green color. *Proteus* and *Providencia* groups give a positive reaction in 1-5 minutes. Other members of *Enterobacteriaceae* give negative reactions.

QUALITY CONTROL

Dehydrated medium

Appearance: free-flowing, homogeneous.

Color: light tan.

Prepared medium

Appearance: slightly opalescent without precipitate.

Color: light amber.

Incubation conditions: 36 ± 1 °C for 18-24 hours.

Microorganism	ATCC	Growth	Reaction
<i>Escherichia coli</i> □ □	25922	good	-
<i>Enterobacter aerogenes</i>	13048	good	-
<i>Proteus mirabilis</i>	25933	good	+
<i>Proteus vulgaris</i>	13315	good	+

STORAGE

The powder is very hygroscopic: store the powder at 10-30 °C, in a dry environment, in its original container tightly closed and use it before the expiry date on the label or until signs of deterioration or contamination are evident.

Store prepared tubes at 2-8 °C.

REFERENCES

1. Ann. Inst. Pasteur. (1954) **87**: 375-386.
2. Pub. Hlth Lab. (1957) **15**: 153.

PRESENTATION

Product	REF	Σ
PHENYLALANINE AGAR (19.2 l)	610039	500 g
PHENYLALANINE AGAR (3.8 l)	620039	100 g

TABLE OF SYMBOLS

LOT Batch code	 Caution, consult accompanying documents	 Manufacturer	 Contains sufficient for <n> tests	 Keep away from heat source
REF Catalogue number	 Fragile, handle with care	 Use by	 Temperature limitation	



LIOFILCHEM s.r.l.

Via Scozia, Zona Ind.le - 64026, Roseto D.A. (TE) - ITALY

Phone +390858930745 Fax +390858930330

Website: www.liofilchem.net E-mail: liofilchem@liofilchem.net